

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/77178 A1

- (51) International Patent Classification⁷: C12N 9/00 (74) Agent: FARRELL, Kevin, M.; P.O. Box 999, York Harbor, ME 03911 (US).
- (21) International Application Number: PCT/US00/16551
- (22) International Filing Date: 15 June 2000 (15.06.2000) (81) Designated States (*national*): AU, CA, JP.
- (25) Filing Language: English (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data: 60/139,408 16 June 1999 (16.06.1999) US
Published:
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- (71) Applicant: BOSTON BIOMEDICAL RESEARCH INSTITUTE [US/US]; 64 Grove Street, Watertown, MA 02472 (US).
- (72) Inventor: RASO, Victor; 69B Strathmore Road, Brighton, MA 02135 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 00/77178 A1

(54) Title: IMMUNOLOGICAL CONTROL OF β -AMYLOID LEVELS *IN VIVO*

(57) Abstract: The present invention provides an antibody which catalyzes hydrolysis of β -amyloid at a predetermined amide linkage. The antibody preferentially binds a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage. Specific antibodies provided include those which catalyze the hydrolysis at the amyloid linkages between residues 39 and 40, 40 and 41, and 41 and 42 of β -amyloid. The present antibody also provides a vectorized antibody which is characterized by the ability to cross the blood brain barrier and also catalyze the hydrolysis of β -amyloid. Also provided are methods for sequestering free β -amyloid in the blood stream, for reducing levels of β -amyloid in the brain, for reducing the level of circulating β -amyloid, for preventing the formation of amyloid plaques in the brain and for disaggregating amyloid plaques. Finally, this invention also provides a method of generating antibodies by immunizing an animal with antigen comprised of an epitope which has a statine analog or which utilizes reduced peptide bond analogs to mimic the conformation of a hydrolysis transition state of a polypeptide.

IMMUNOLOGICAL CONTROL OF β -AMYLOID LEVELS IN VIVOBackground of the Invention

Alzheimer's disease is a progressive and ultimately fatal form of dementia that affects a substantial portion of the elderly population. Definitive diagnosis at autopsy relies on the presence of neuropathological brain lesions marked by a high density of senile plaques. These extracellular deposits are found in the neo-cortex, hippocampus and amygdala as well as in the walls of the meningeal and cerebral blood vessels. The principal component of these plaques is a 39 to 43 residue β -amyloid peptide. Each plaque contains approximately 20 fmole (80 picograms) of this 4 kDa peptide (Selkoe et al., *J. of Neurochemistry* 46: 1820 (1986)). Apolipoprotein E and neurofibrillary tangles formed by the microtubule-associated tau protein are also often associated with Alzheimer's disease.

β -amyloid is proteolytically cleaved from an integral membrane protein called the β -amyloid precursor protein. The gene which codes for this protein in humans is found on chromosome 21 (St George-Hyslop et al., *Science* 235: 885 (1987), Kang et al., *Nature* 325: 733 (1987)). Numerous cultured cells and tissues (eg. brain, heart, spleen, kidney and muscle) express this β -amyloid precursor protein and also secrete the 4 kDa β -amyloid fragment into culture media, apparently as part of a normal processing pathway.

While it is difficult to establish an absolute causal relationship between β -amyloid or the plaques it forms and Alzheimer's disease, there is ample evidence to support the pathogenic role of β -amyloid. For example, patients with Down's syndrome have an extra copy of the β -amyloid precursor protein gene due to trisomy of chromosome 21 (St George-Hyslop et al., *Science* 235: 885 (1987), Kang et al., *Nature* 325: 733 (1987)). They correspondingly develop an early-onset Alzheimer's disease neuropathology at 30-40 years of age. Moreover, early-onset familial Alzheimer's disease can result from

mutations in the β -amyloid precursor protein gene which fall within or adjacent to the β -amyloid sequence (Hardy, J., *Nature Genetics* 1: 233 (1992)). These observations are consistent with the notion that deposition of β -amyloid as plaques in the brain are accelerated by an elevation in its extracellular concentration (Scheuner et al., *Nature Med.* 2: 864 (1996)). The finding that β -amyloid is directly neurotoxic both in vitro and in vivo (Kowall et al., *Proc. Natl. Acad. Sci.* 88: 7247 (1991)), suggest that soluble aggregated β -amyloid, not the plaques per se, may produce the pathology.

Observations have indicated that amyloid plaque formation may proceed by a crystallization type mechanism (Jarrett et al., *Cell* 73: 1055 (1993)). According to this model, the seed that initiates plaque nucleation is an β -amyloid which is 42 or 43 amino acids long ($A\beta_{1-43}$). The rate-determining nucleus formed by $A\beta_{1-43}$ or $A\beta_{1-42}$ allows peptides $A\beta_{1-40}$ or shorter to contribute to the rapid growth of an amyloid deposit. This nucleation phenomenon was demonstrated in vitro by the ability of $A\beta_{1-42}$ to cause the instantaneous aggregation of a kinetically stable, supersaturated solution of $A\beta_{1-40}$. That finding has led to the possibility that $A\beta_{1-40}$ might be relatively harmless in the absence of the nucleation peptides $A\beta_{1-42}$ or $A\beta_{1-43}$. Indeed, elevated levels of these long peptides have been found in the blood of patients with familial Alzheimer's disease (Scheuner et al., *Nature Med.* 2: 864 (1996)). Moreover, $A\beta_{1-42}$ or $A\beta_{1-43}$ was found to be the predominant form deposited in the brain plaques of many Alzheimer's disease patients (Gravina et al., *J. of Biol. Chem.* 270: 7013 (1995)).

Given the central role played by β -amyloid, it has become increasingly important to understand the interrelationship between the different pools of these molecules in the body. Free β -amyloid present in the blood most likely arises from peptide released by

-3-

proteolytic cleavage of β -amyloid precursor protein present on cells in the peripheral tissues. Likewise most of the free β -amyloid found in the brain and cerebrospinal fluid is probably derived from peptide released by secretase cleavage of β -amyloid precursor protein expressed on brain cells. The peptides are identical regardless of origin, and the results from several studies suggest an intercommunication between these pools.

10 Summary of the Invention

One aspect of the present invention is an antibody which catalyzes hydrolysis of β -amyloid at a predetermined amide linkage. In one embodiment, the antibody preferentially binds a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage and also binds to natural β -amyloid with sufficient affinity to detect using an ELISA. In another embodiment, the antibody preferentially binds a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage, and does not bind natural β -amyloid with sufficient affinity to detect using an ELISA. Antibodies generated are characterized by the amide linkage which they hydrolyze. Specific antibodies include those which catalyze the hydrolysis at the amyloid linkages between residues 39 and 40, 40 and 41, and 41 and 42, of β -amyloid.

Another aspect of the present invention is a vectorized antibody which is characterized by the ability to cross the blood brain barrier and is also characterized by the ability to catalyze the hydrolysis of β -amyloid at a predetermined amide linkage. In one embodiment, the vectorized antibody is a bispecific antibody. Preferably, the vectorized antibody has a first specificity for the transferrin receptor and a second specificity for a transition state adopted by β -

amyloid during hydrolysis. Specific vectorized antibodies include those which catalyze the hydrolysis at the amyloid linkages between residues 39 and 40, 40 and 41, and 41 and 42, of β -amyloid.

- 5 Another aspect of the present invention is a method for sequestering free β -amyloid in the bloodstream of an animal by intravenously administering antibodies specific for β -amyloid to the animal in an amount sufficient to increase retention of β -amyloid in the circulation.
- 10 Therapeutic applications of this method include treating patients diagnosed with, or at risk for Alzheimer's disease.

- Another aspect of the present invention is a method for sequestering free β -amyloid in the bloodstream of an
- 15 animal by immunizing an animal with an antigen comprised of an epitope which is present on β -amyloid endogenous to the animal under conditions appropriate for the generation of antibodies which bind endogenous β -amyloid. Therapeutic applications of this method include treating
- 20 patients diagnosed with, or at risk for Alzheimer's disease.

- Another aspect of the present invention is a method for reducing levels of β -amyloid in the brain of an animal by intravenously administering antibodies specific
- 25 for endogenous β -amyloid to the animal in an amount sufficient to increase retention of β -amyloid in the circulation of the animal. In one embodiment, the antibodies are catalytic antibodies which catalyze hydrolysis of β -amyloid at a predetermined amide linkage.
- 30 The antibodies may be either monoclonal or polyclonal. In one embodiment, the antibodies specifically recognize epitopes on the C-terminus of β -amyloid₁₋₄₃.

- Another aspect of the present invention is a method for reducing levels of β -amyloid in the brain of an
- 35 animal, by immunizing the animal with an antigen comprised of an epitope which is present on endogenous β -amyloid under conditions appropriate for the generation

of antibodies which bind endogenous β -amyloid. In one embodiment, the antigen is a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage. In a preferred embodiment, the antigen is comprised of $A\beta_{10-25}$. Preferably, the antibodies generated have a higher affinity for the transition state analog than for natural β -amyloid, and catalyze hydrolysis of endogenous β -amyloid.

Similar methods which utilize or generate antibodies which catalyze the hydrolysis of β -amyloid for reducing levels of circulating β -amyloid in an animal, and also for preventing the formation of amyloid plaques in the brain of an animal, are also provided. Also, methods for disaggregating amyloid plaques present in the brain of an animal by utilizing or generating antibodies which catalyze the hydrolysis of β -amyloid are provided.

Another aspect of the present invention is a method for disaggregating amyloid plaques present in the brain of an animal by intravenously administering vectorized bispecific antibodies to the animal in an amount sufficient to cause significant reduction in β -amyloid levels in the brain of the animal. The vectorized bispecific antibodies are competent to transcytose across the blood brain barrier, and have the ability to catalyze hydrolysis of endogenous β -amyloid at a predetermined amide linkage upon binding. Preferably, the vectorized bispecific antibodies specifically bind the transferrin receptor.

Another aspect of the present invention is a method for generating antibodies which catalyze hydrolysis of a protein or polypeptide by immunizing an animal with an antigen comprised of an epitope which has a statine analog which mimics the conformation of a predetermined hydrolysis transition state of the polypeptide, under conditions appropriate for the generation of antibodies to the hydrolysis transition state. This method can be

used to generate catalytic antibodies to β -amyloid. A similar method, which utilizes reduced peptide bond analogs to mimic the conformation of a hydrolysis transition state of a polypeptide, is also provided.

5 Brief Description of the Figures

Figure 1 is an amino acid sequence listing (SEQ ID NO: 1) of the 43 residue β -amyloid peptide ($A\beta$).

Figure 2 is an amino acid sequence listing (SEQ ID NO: 2) of the antigenic peptide made from the N-terminal
10 sequence of β -amyloid ($A\beta_{1-16}$).

Figure 3 is an amino acid sequence listing (SEQ ID NO: 3) of the antigenic peptide made from the central region of β -amyloid ($A\beta_{10-25}$).

Figure 4 is an amino acid sequence listing (SEQ ID
15 NO: 4) ($A\beta_{35-43}$) of the antigenic peptide made from the C-terminal sequence of β -amyloid.

Figure 5 is a diagrammatic representation of data from an ELISA comparing monoclonal antibody binding to $A\beta_{35-43}$ and $A\beta_{1-43}$ versus $A\beta_{1-40}$.

20 Figure 6 indicates the amide linkages in the peptide made from the β -amyloid C-terminal sequence (SEQ ID NO: 4) that were independently replaced with a statyl moiety, to generate the different statine transition state analogs of the peptide.

25 Figure 7 indicates the amide linkages in the peptide made from the β -amyloid central sequence (SEQ ID NO: 3) that were independently replaced with a statyl moiety, to generate the different phenylalanine statine transition state analogs of the peptide.

30 Figure 8 is a structural comparison between the native β -amyloid peptide and the transition state phenylalanine statine β -amyloid peptide analog.

Figure 9 is a structural comparison between the native β -amyloid peptide and the reduced peptide bond
35 transition state β -amyloid peptide analog.

Figure 10 is a formulaic representation of the native C-terminal region of β -amyloid, and the phosphoramidate transition state analog of the C-terminal region of β -amyloid ($A\beta_{35-43}$).

5 Figure 11 indicates the putative transition state for peptide hydrolysis by zinc peptidases, compared to the phosphonate and phosphoramidate mimics.

Figure 12 is a structural comparison of the native β -amyloid peptide and the transition state
10 phosphoramidate β -amyloid peptide which has the peptide link between Gly 38 and Val 39 replaced with a phosphoramidate bond.

Figure 13 is a diagrammatic representation of data from an ELISA which assess the binding of monoclonal
15 antibodies generated to transition state β -amyloid peptide analogs, to the normal $A\beta_{1-43}$ and to the phenylalanine statine transition state β -amyloid peptide.

Figure 14 is a diagrammatic representation of data from an ELISA comparing antibody binding to the statine
20 transition state β -amyloid peptide versus native $A\beta_{1-43}$ and native $A\beta_{1-40}$.

Figure 15 is a graph of data showing the cleavage of ^{125}I - $A\beta$ -sepharose by monoclonal antibodies generated to transition state analogs of β -amyloid.

25 Figure 16 is a diagrammatic representation of data which quantitate the attachment of bispecific antibody to receptor-positive cells.

Figure 17 is a diagrammatic representation of data obtained from experiments designed to track the
30 transcytosis of vectorized bispecific antibody into brain.

Detailed Description of the Invention

The present invention relates to immunologically based methods for controlling levels of β -amyloid in the
35 body of an animal. The invention is based on the finding that antibodies specific for β -amyloid are able to bind

β -amyloid in the presence of a physiological level of human serum albumin. The invention is also based on the finding that an animal can tolerate the presence of antibodies specific for β -amyloid in amounts sufficient to sequester β -amyloid in the bloodstream.

One aspect of the present invention relates to a method for sequestering free β -amyloid in the bloodstream of an animal. The soluble and insoluble forms of β -amyloid present within an animal are in dynamic equilibrium. Soluble β -amyloid is thought to translocate between blood and cerebrospinal fluid. Insoluble β -amyloid aggregates deposit from the soluble pool in the brain, as amyloid plaques. Results detailed in the Exemplification section below indicate that intravenous administration of antibodies specific for β -amyloid to an animal impedes the passage of soluble β -amyloid out of the peripheral circulation. This occurs because the β -amyloid specific antibodies, which are restricted to the peripheral circulation, bind to β -amyloid and sequester it in the circulation. Such sequestration is accomplished through intravenous administration of an appropriate amount of antibodies specific for β -amyloid to the animal. The amount of antibody which is sufficient to produce sequestration is dependent upon various factors (e.g., specific characteristics of the antibody to be delivered, the size, metabolism, and overall health of the animal) and are to be determined on a case by case basis.

Administered antibodies can be monoclonal antibodies, a mixture of different monoclonal antibodies, polyclonal antibodies, or any combination therein. In one embodiment, the antibodies bind to the C-terminal region of β -amyloid. Such antibodies specifically bind the less abundant, but more noxious $A\beta_{1-43}$ species in the blood as opposed to the smaller and less detrimental $A\beta_{1-40}$. In another embodiment, a combination of antibodies having specificity for various regions of β -amyloid are

administered. In another embodiment, antibodies which catalyze the hydrolysis of β -amyloid, discussed in more detail below, are administered either alone or in combination with other anti- β -amyloid antibodies.

5 The animal to which the antibodies are administered is any animal which has circulating soluble β -amyloid. In one embodiment, the animal is a human. The human may be a healthy individual, or alternatively, may be suffering from or at risk for a disease in which elevated
10 β -amyloid levels are thought to play a role, for example a neurodegenerative disease such as Alzheimer's disease.

A related aspect of the present invention is a method for sequestering free β -amyloid in the bloodstream of an animal by stimulating an immune response within the
15 animal to endogenous β -amyloid. The results detailed in the Exemplification below indicate that an animal can tolerate the induction of an immune response which produces antibodies to endogenous β -amyloid, and that the presence of such antibodies will alter the distribution
20 of β -amyloid in the body, in a similar manner as the above described method of administering β -amyloid binding antibodies. The immune response to endogenous β -amyloid is generated by immunizing the animal with one or more antigens comprised of epitopes present on the endogenous
25 β -amyloid. Epitopes present on the inoculated antigens can correspond to epitopes present on any region of the β -amyloid molecule. In a preferred embodiment, epitopes found on the C-terminal region of β -amyloid are used to generate antibodies which specifically bind the $A\beta_{1-43}$
30 species as opposed to the smaller $A\beta_{1-40}$. In an alternate embodiment, a combination of different epitopes are administered to generate a variety of antibodies to β -amyloid. A more generalized immune response is generated by immunizing either with a mixture of different small
35 peptide antigens or with the full-length 43 residue β -amyloid peptide. In another embodiment, antigens used for inoculation include transition state analogs of β -

-10-

amyloid peptides to induce antibodies which have catalytic activity directed towards β -amyloid hydrolysis, described in detail below.

5 The immunoreactivity of the antigens can be enhanced by a variety of methods, many of which involve coupling the antigen to an immunogenic carrier. In addition, various methods are known and available to one of skill in the art for specifically enhancing the immunogenicity of endogenous molecules or the epitopes contained
10 therein. Various modifications can be made to the β -amyloid antigen(s) described herein to render it more compatible for human use. For example, the peptide(s), can be genetically engineered into appropriate antigenic carriers, or DNA vaccines can be designed.

15 The above techniques for sequestering β -amyloid in the circulation are also useful for reducing the levels of β -amyloid in the brain. Because the formation of amyloid plaques in the brain is dependent, at least in part, on the levels of free β -amyloid present in the
20 brain, reducing brain β -amyloid levels of an animal will, in turn, reduce the formation of amyloid plaques in the brain. Therefore, the above techniques are useful for preventing the formation of amyloid plaques in the brain of an animal. This is especially applicable to an animal
25 which is considered at risk for the development of amyloid plaques; a risk which may result from a genetic predisposition or from environmental factors.

Administration of antibodies, or immunization of the animal to produce endogenous antibodies, to β -amyloid can
30 be of therapeutic benefit to such an animal (e.g., a human who has a family history of Alzheimer's disease, or who is diagnosed with the disease).

Another aspect of the present invention relates to antibodies which are characterized by the ability to
35 catalyze the hydrolysis of β -amyloid at a predetermined amide linkage. Experiments detailed in the Exemplification section demonstrate the generation of

different antibodies which have proteolytic activity towards β -amyloid. Such antibodies are generated by immunizing an animal with an antigen which is a transition state analog of the β -amyloid peptide. A transition state analog mimics the transition state that β -amyloid adopts during hydrolysis of a predetermined amide linkage. Transition state analogs useful for generating the catalytic antibodies include, without limitation, statine, phenylalanine statine, phosphonate, phosphoramidate, and reduced peptide bond transition state analogs.

Antibodies generated to epitopes unique to the transition state preferentially bind β -amyloid in the transition state. Binding of these antibodies stabilizes the transition state, which leads to hydrolysis of the corresponding amide bond. The particular amide linkage to be hydrolyzed is chosen based upon the desired cleavage product. For example, cleavage of full length β -amyloid into two peptide fragments which cannot aggregate into amyloid plaques would be of therapeutic use in the methods disclosed herein. Antibodies may be either monoclonal or polyclonal. Several such transition state mimics have been made and used as antigen in the generation of monoclonal antibodies which catalyze the cleavage at the indicated linkage. These antigens and the antibodies generated are listed in Table 8 of the Exemplification section below. Antibodies generated to antigens which have transition state mimics incorporated at a specific amide linkage, should bind the natural hydrolysis transition states of these linkages in native β -amyloid, stabilizing the transition state and catalyzing cleavage at that linkage.

At least two different classes of antibodies are generated by the above methods. The first class preferentially binds the transition state analog, and also detectably cross reacts with natural β -amyloid using the ELISA detailed in the Exemplification section, to

-12-

detect binding. The second class binds the transition state analog, and does not detectably cross react with natural β -amyloid using the ELISA procedure detailed in the Exemplification section to detect binding. Both

5 classes of antibodies have potential value as catalytic antibodies. The respective binding affinities of an anti-transition state antibody is likely to reflect its activity at catalyzing hydrolysis. It is thought that in order for an antibody to have activity at catalyzing

10 hydrolysis of a protein, it must possess at least a minimal ability to bind the natural (non-transition) state of the protein. Antibodies which retain significant binding for β -amyloid, (that strongly cross react with natural β -amyloid) may be more efficient at

15 catalyzing hydrolysis due to a higher efficiency of binding the β -amyloid. Once bound, these antibodies force the protein into a transition state conformation for hydrolytic cleavage. Alternatively, antibodies which only minimally cross react with natural β -amyloid,

20 although less efficient at binding native β -amyloid, are likely to be more efficient at forcing the bound β -amyloid into the transition state conformation for hydrolytic cleavage. It should be pointed out that failure to detect binding of the anti-transition state

25 antibodies to natural β -amyloid by the ELISA methods presented in the Exemplification herein does not necessarily reflect an inability to bind natural β -amyloid sufficiently to function as a catalytic antibody. More likely, a lack of detection merely reflects the

30 sensitivity limitations of the assay.

Antibodies which have substantial affinity for the predicted cleavage products of the native β -amyloid peptide may be subject to product inhibition and might therefore exhibit low turnover. Such undesirable

35 antibodies can be identified by secondary screening using peptides which contain epitopes of the predicted cleavage products (e.g., via ELISA).

In a preferred embodiment, the antibodies are monoclonal. Monoclonal antibodies are produced by immunizing an animal (e.g., mouse, guinea pig, or rat) with the transition state analog antigen, and subsequently producing hybridomas from the animal, by standard procedures. Hybridomas which produce the desired monoclonal antibodies are identified by screening. One example of a screening method is presented in the Exemplification section which follows.

In another embodiment, the antibodies are polyclonal. Polyclonal antibodies are generated by immunizing an animal (e.g., a rabbit, chicken, or goat) with antigen and obtaining sera from the animal. Polyclonal antibodies which have the desired binding specificities can be further purified from the sera by one of skill in the art through the course of routine experimentation.

Catalytic antibodies specific for β -amyloid can alternatively be generated in an individual through the use of anti-idiotypic vaccines designed to elicit the production of catalytic antibodies. Such vaccines are described in the disclosure of Raso and Paulus (U.S. Patent Application 09/102,451, ANTI-IDIOTYPE VACCINES TO ELICIT CATALYTIC ANTIBODIES, filed by Applicants June 22, 1998, currently pending), the contents of which are incorporated herein by reference.

Another aspect of the present invention is the use of statine and reduced peptide bond analogs to elicit catalytic antibodies having proteolytic activity. The Exemplification section below details methods for using statine analogs as antigen in the production of catalytic antibodies, and also lists examples of anti-transition-state antibodies generated using these methods. The "statyl" moiety is derived from naturally evolved protease transition state inhibitors like amastatin, pepstatin, and bestatin. These naturally-occurring statine-based inhibitors have been used to effectively block the activity of aminopeptidases, aspartic proteases

and the HIV protease. Synthetic peptides containing a statine residue offer novel features for the induction of catalytic antibodies. The statyl moiety has a tetrahedral bond geometry, its length is extended by two
5 CH₂ units, it has a strategically placed OH group and the structure has no charge. The presence of the additional CH₂ units is expected to elicit a more elongated antibody combining site, and antibodies possessing this extended site will induce extra strain on the peptide substrate,
10 producing an accelerated catalysis. In addition, the -OH group in these statine analogs is thought to better approximate the position and chemistry of the true transition state. Statine-based transition-state analogs should therefore elicit a class of antibodies
15 which is significantly different from those obtained from the more commonly used negatively charged phosphonate analogs.

Reduced peptide bond analogs introduce a tetrahedral configuration, without increasing the distance between
20 amino acid residues. This feature should more closely approximate the true transition state geometry, than previously used analogs. A positively charged secondary amine replaces the amide nitrogen of the natural polypeptide and should elicit a complementary negatively
25 charged side chain at a proximal locus in the antibody combining site. The presence of such ancillary glutamyl or aspartyl groups on the antibody will assist antibody-mediated catalysis of peptide cleavage via acid-base exchange. Reduced peptide bond-based transition-state
30 analogs should therefore elicit a class of antibodies which is significantly different from those obtained from using the more commonly used negatively charged phosphonate analogs. Reduced peptide bond analogs and
statine analogs can be used to produce specific
35 transition state analog antigens for a wide variety of proteins or polypeptides. These antigens can in turn be used to generate the respective catalytic antibodies.

Administration of the β -amyloid catalytic antibodies described above can be used in the methods described above for 1) sequestering free β -amyloid in the bloodstream of an animal, 2) reducing levels of β -amyloid in the brain of an animal, and 3) preventing the formation of amyloid plaques in the brain of an animal, to generate the analogous results. Experiments presented in the Exemplification demonstrate that immunization of an animal with a transition state analog results in the generation of an immune response to produce antibodies which recognize the transition state, and which catalyze hydrolysis of the β -amyloid protein. This indicates that the transition state analogs can be used as antigens in these methods to induce the production of antibodies in the animal which recognize and catalyze cleavage of endogenous β -amyloid.

Methods which involve reducing overall levels of β -amyloid in an animal through the proteolytic action of the above described catalytic antibodies are also encompassed by the present invention. The presence of functional catalytic antibodies in the circulation of an animal will reduce the level of intact β -amyloid in the circulation by selective hydrolytic cleavage. Accordingly, the present invention provides a method for reducing levels of circulating β -amyloid in an animal by introducing the above described catalytic antibodies into the animal. Administration of the antibodies to the animal is preferably via intravenous administration. Such antibodies are either monoclonal, mixed monoclonal, polyclonal or any mixture thereof. The origin of the antibody may affect the half-life of the antibody in the animal; antibodies from less related species are more likely to be recognized as foreign by the animal's immune system. Preferably, administered antibodies are derived from a species closely related to the animal, to maximize half-life and minimize adverse reactions by the host.

Administration of isolated variable region antibody fragments may produce beneficial results in this regard.

The present invention also provides a method for reducing levels of circulating β -amyloid in an animal by immunizing the animal with a β -amyloid transition state analog to induce endogenous catalytic antibody production. The use and design of such vaccines is described above, and detailed in the Exemplification section below.

10 The reduction of β -amyloid levels in the circulation of an animal is expected to displace the equilibrium of β -amyloid in the body, and ultimately lead to a reduction in the levels of β -amyloid in the brain of the animal through mass action. In this respect, the present
15 invention provides methods for reducing the levels of β -amyloid in the brain of an animal, by either administering catalytic antibodies to the animal, or by administering a transition state analog to induce endogenous antibody production. It follows that these
20 procedures also have value as methods for preventing the formation of amyloid plaques in the brain of an animal, since the resulting reduction in the levels of β -amyloid in the brain of an animal should prevent the formation of amyloid plaques. These procedures also have value as
25 methods for disaggregating amyloid plaques present in the brain of an animal, since evidence indicates that lower brain β -amyloid levels can lead to the disaggregation of plaques.

Another aspect of the present invention provides a
30 more direct method of altering the distribution of β -amyloid in the brain by actually delivering anti- β -amyloid antibodies to the brain. Methods described above for reducing levels of β -amyloid in the brain and for preventing aggregation of amyloid plaques depend upon
35 exchange between β -amyloid pools in the blood, tissues, cerebrospinal fluid and the brain, the exchange being driven by an antibody-mediated disruption of the

equilibrium between these different pools. In contrast, delivery of anti- β -amyloid antibodies to the brain will directly affect β -amyloid aggregation. Evidence presented in the Exemplification section below indicates that the binding of certain anti- β -amyloid antibodies inhibits the initial aggregation of β -amyloid *in vitro*, and also disaggregates preformed *in vitro* β -amyloid complexes. Moreover, if insoluble peptide is in equilibrium with a low level of soluble β -amyloid, then an anti- β -amyloid binding antibody could upset this balance and gradually dissolve the precipitate. These observations indicate that the presence of β -amyloid antibodies in the brain will directly inhibit the formation of amyloid plaques and will also disaggregate preformed plaques by disrupting the dynamic equilibrium between soluble β -amyloid and fibrillar β -amyloid deposited as plaques. Furthermore, a highly active catalytic antibody is expected to destroy insoluble β -amyloid plaques by hydrolytically cleaving the constituent aggregated peptides.

One way of delivering antibodies to the brain is by producing vectorized antibodies competent for transcytosis across the blood-brain barrier. Vectorized antibodies are produced by covalently linking an antibody to an agent which promotes delivery from the circulation to a predetermined destination in the body. Examples of vectorized molecules which can traverse the blood-brain barrier are found in the prior art (Bickel et al., *Proc. Natl. Acad. Sci. USA* 90: 2618-2622 (1993); Broadwell et al., *Exp. Neurol.* 142: 47-65 (1996)). In these examples, antibodies are linked to another macromolecule, the antibodies being the agent which promotes delivery of the macromolecules. One example of such an agent is an antibody which is directed towards a cell surface component, such as a receptor, which is transported away from the cell surface. Examples of antibodies which confer the ability to transcytose the blood-brain barrier

include, without limitation, anti-insulin receptor antibodies, and also anti-transferrin receptors (Saito et al., *Proc. Natl. Acad. Sci. USA* 92: 10227-31 (1995); Pardridge et al., *Pharm. Res.* 12: 807-816 (1995); Broadwell et al., *Exp. Neurol.* 142: 47-65 (1996)). This first antibody is covalently linked to an antibody which binds β -amyloid. Alternatively, coupling the β -amyloid antibodies to ligands which bind these receptors (e.g., insulin, transferrin, or low density lipoprotein) will also produce a vectorized antibody competent for delivery to the brain from the circulation (Descamps et al., *Am. J. Physiol.* 270: H1149-H1158 (1996); Duffy et al., *Brain Res.* 420: 32-38 (1987); Dehouck et al., *J. Cell Biol.* 138: 877-889 (1997)).

A vector moiety can be chemically attached to the anti- β -amyloid antibody to facilitate its delivery into the central nervous system. Alternatively, the moiety can be genetically engineered into the antibody as an integral component. This vector component can be for example, an anti-transferrin receptor antibody or anti-insulin receptor antibody which binds the receptors present on the brain capillary endothelial cells (Bickel et al., *Proc. Natl. Acad. Sci. USA* 90: 2618-22 (1993); Pardridge et al., *J. Pharmacol. Exp. Ther.* 259: 66-70 (1991); Saito et al., *Proc. Natl. Acad. Sci. USA* 92: 10227-31 (1995); Friden et al., *J. Pharm. Exper. Ther.* 278: 1491-1498 (1996)) which make up the blood-brain barrier. The resulting bifunctional antibody will attach to the appropriate receptors on the luminal side of the vessel (Raso et al., *J. Biol. Chem.* 272: 27623-27628 (1997); Raso et al., *J. Biol. Chem.* 272: 27618-27622 (1997); Raso, V. *Anal. Biochem.* 222: 297-304 (1994); Raso et al., *Cancer Res.* 41: 2073-2078 (1981); Raso et al., Monoclonal antibodies as cell targeted carriers of covalently and non-covalently attached toxins. In Receptor mediated targeting of drugs, vol. 82. G. Gregoriadis, G. Post, J. Senior and A. Trouet, editors.

NATO Advanced Studies Inst., New York. 119-138 (1984)).
Once bound to the receptor, both components of the
bisppecific antibody pass across the blood-brain barrier
by the process of transcytosis. Anti- β -amyloid
5 antibodies which have entered the brain interact directly
with both β -amyloid plaques and the soluble β -amyloid
pool. It has been estimated that concentrations of
macromolecules in the 10^{-8} - 10^{-7} M range can be achieved in
the brain using vector-mediated delivery via these brain
10 capillary enriched protein target sites (Maness et al.,
Life Sciences 55: 1643-1650 (1994); Lerner et al.,
Science 252: 659-667 (1991)). Importantly, the vector
appears safe since animals dosed daily for two weeks with
an anti-transferrin receptor antibody displayed no loss
15 of integrity of the blood-brain barrier, using a
radioactive sucrose probe (Broadwell et al., *Exp. Neurol.*
142: 47-65 (1996)).

The Exemplification details the production of
vectorized bisppecific antibodies which bind β -amyloid.
20 The bisppecific antibodies transcytose across the blood
brain barrier via a first specificity which binds the
transferrin receptor. Use of antibodies which bind the
transferrin receptor for delivery of agents across the
blood brain barrier is described by Friden et al. in U.S.
25 Patent No. 5,182,107; No. 5,154,924; No. 5,833,988; and
No. 5,527,527; the contents of which are incorporated
herein by reference.

Results from experiments presented in the
Exemplification section which follows indicate that the
30 produced bisppecific antibodies retain their separate
specificities and are delivered across the blood-brain
barrier into the brain parenchyma and brain capillaries
of a live animal when administered intravenously.

Alternate methods for the production of bisppecific
35 antibodies have been described for genetically
engineering bisppecific reagents or for producing them
intracellularly by fusing the two different hybridoma

-20-

clones (Holliger et al., *Proc. Natl. Acad. Sci.* 90: 6444-6448 (1993); Milstein et al., *Nature* 305: 537 (1983); Mallander et al., *J. Biol. Chem.* 269: 199-206 (1994)).

Vectorized bispecific antibodies produced by these
5 techniques can also be used in the methods of the present invention.

Since the introduction of whole antibodies into the brain might be detrimental if they were to fix complement and promote complement-mediated lysis of neuronal cells,
10 it may be beneficial to produce and utilize smaller vectorized $F(ab')_2$ bispecific reagents. It has been shown that aggregated β -amyloid itself can fix complement in the absence of any antibody and that the resulting inflammation may contribute to the pathology of
15 Alzheimer's disease. The possibility of intracerebral antibody having a similar effect can be greatly reduced by eliminating the Fc region of the antibody. Moreover, since coupling of Fab' halves uses the intrinsic hinge region cysteines, no extraneous substituent linkage
20 groups need be added. Faster or more efficient entry into the brain represents another potential advantage that smaller $F(ab')_2$ or Fv_2 reagents may provide for intracerebral delivery. In addition, the two types of vectorized molecules may have different biodistribution
25 and plasma half-life characteristics (Spiegelberg et al., *J. Exp. Med.* 121: 323 (1965)).

Depending on their design, anti- β -amyloid bispecific antibodies in the brain can reduce soluble β -amyloid and β -amyloid deposits by three potential mechanisms. An
30 anti- β -amyloid bispecific antibody that tightly binds soluble β -amyloid will not only sequester the peptide but, due to efflux of vectorized molecules from the central nervous system (Kang et al., *J. Pharm. Exp. Ther.* 269: 344-350 (1994)), may also carry the bound β -amyloid
35 out of the brain, releasing it into the blood stream. Such a clearance mechanism would lead to a continuous cycling of β -amyloid out of the brain. In addition, if

the antibodies have catalytic activity, they will directly reduce the levels of harmful β -amyloid by degradation. Since catalytic antibodies exhibit turnover, each antibody can inactivate many β -amyloid molecules. Thus much less vectorized bispecific antibody has to be delivered into the brain to achieve the desired depletion of β -amyloid.

To be effective the anti- β -amyloid sites of a bispecific antibody must be empty before passage out of the blood and into the brain. Therefore the concentration of bispecific antibody in animals must exceed the level of β -amyloid circulating in the blood. Calculations performed based upon known β -amyloid levels (Scheuner et al., *Nature Med.* 2: 864-870 (1996)) and a medium-range plasma level of bispecific antibody expected in a treated animal indicated 99.9% of the bispecific antibodies that enter the brain will have unoccupied anti- β -amyloid combining sites.

Another way of delivering antibodies to the brain is via direct infusion of anti- β -amyloid antibodies into the brain of an animal. This technique gives these antibodies immediate access to β -amyloid in the brain without having to cross the blood-brain barrier. Direct infusion can be accomplished via direct parenchymal or intracerebroventricular infusion (Knopf et al., *J. Immunol.* 161: 692-701 (1998)). Briefly, the animal is anesthetized and placed in a stereotaxic frame. A midsagittal incision is made on the scalp to expose the skull and the underlying fascia is scraped away. A hole is drilled to accept a sterilized length of stainless steel hypodermic tubing, which is stereotaxically advanced so that its tip is appropriately located in the brain. A guide cannula is then attached to the skull and sealed. The cannula remains in place for multiple infusions of antibody into the brain. A bolus of a sterile 50 mg/ml solution of a monoclonal anti- β -amyloid

-22-

can be infused over a 2-8 minute period into an immobilized animal via an injection cannula.

Delivery of catalytic antibodies into the brain of an animal via one of the above described methods, can also be used to disaggregate amyloid plaques present in the brain. The advantage of delivering an β -amyloid-specific catalytic antibody into the brain is two-fold. The β -amyloid peptide is permanently destroyed by such antibodies and, since catalysis is continuous, each antibody inactivates many target β -amyloid molecules in the brain. Thus much less antibody has to be infused into the central nervous system to achieve the desired depletion of β -amyloid.

The amount of antibody to be administered or delivered to the animal should be sufficient to cause a significant reduction in β -amyloid levels in the brain of the animal. The appropriate amount will depend upon various parameters (e.g., the particular antibody used, the size and metabolism of the animal, and the levels of endogenous β -amyloid) and is to be determined on a case by case basis. Such determination is within the means of one of average skill in the art through no more than routine experimentation.

It is expected that additional benefits with respect to lowering brain β -amyloid levels and preventing or disaggregating amyloid plaques can be achieved through utilizing a combination of one or more of the above described approaches.

Exemplification

30 SECTION 1: RETENTION OF β -AMYLOID IN THE CIRCULATION

Synthesis of β -Amyloid Peptide Antigens

The amino acid sequence of the 43 residue β -amyloid peptide ($A\beta$) is listed in Figure 1. To determine which sites on this $A\beta$ peptide were best suited for antibody-

mediated therapy, three key regions (amino-terminal, central and carboxy-terminal) of the A β 43-mer were chosen to generate epitope-specific vaccines. These shortened peptides served as antigenic epitopes to induce
5 a highly specific antibody response.

Monoclonal antibodies to the amino-terminal region of A β have been shown in the past to have the ability to solubilize A β aggregates (Solomon et al., *Proc. Natl. Acad. Sci. USA* 94(8): 4109 (1997); Solomon et al., *Proc. Natl. Acad. Sci. USA* 93(1): 452 (1996)). A peptide
10 consisting of the amino-terminal region of A β was similarly designed for the present experiments (shown in Fig. 2 and listed in SEQ ID NO: 2) and used to elicit amino-terminal specific antibodies that bind A β . A Cys
15 residue was added to the C-terminus of the A β sequence to provide a suitable linkage group for coupling this peptide to an antigenic carrier protein such as maleimide-activated Keyhole Limpet Hemocyanin (KLH).

A peptide encompassing the central region of A β was
20 also synthesized (shown in Figure 3 and listed in SEQ ID NO: 3). A Cys residue was placed at the N-terminus of the A β sequence to provide a sulfhydryl linkage group for coupling the peptide to antigenic carrier proteins such as maleimide-activated KLH.

To produce an antigen for eliciting an immune
25 response directed against the carboxy-terminus of A β (Suzuki et al., *Science* 264: 1336 (1994)), a decapeptide encompassing the N-terminal region of A β , with an additional Cys residue at the N-terminus, was synthesized
30 (Shown in Fig. 4, and listed in SEQ ID NO: 4). The Cys substitution was designed to provide a sulfhydryl linkage group for coupling the peptide to antigenic carrier proteins such as KLH.

Coupling the Peptides to an Antigenic Carrier Protein

35 The different Cys containing A β peptides were individually thioether-linked to maleimide-activated KLH.

-24-

A multivalent A β vaccine was also produced by simultaneously linking all three of these peptides to maleimide-activated KLH. In addition the full-length A β 43-mer was linked to KLH using glutaraldehyde.

5 Antibodies Elicited with the β -Amyloid Vaccines

Normal BALB/c mice were immunized by standard procedures with the KLH-linked A β vaccines described above. The mice were either bled or sacrificed for removal of the spleen for hybridoma production. Sera and
10 monoclonal antibodies obtained were characterized for binding to A β .

Table 1 shows the results from an ELISA run with 1/100 diluted serum from two non-immunized control mice versus 1/100 and 1/1000 diluted serum from a mouse that
15 was immunized with a central region A β peptide-KLH vaccine. The free A β peptide was adsorbed directly onto the microtitre plate to avoid detection of anti-KLH antibodies in the serum.

Table 1 ELISA for Binding to the Central Region A β Peptide

Bound		Antibody
Addition		(O.D. 450
<u>nm)</u>		
Control Serum A	1/100	0.666
Control Serum B	1/100	0.527
Mouse 1 antiserum	1/100	3.465
Mouse 1 antiserum	1/1000	2.764

Monoclonal antibodies raised against this central region A β peptide and produced by hybridoma fusion were identified using the above described ELISA assay. A binding assay was performed to determine whether the

monoclonal anti-A β antibodies identified also bound to the full length A β peptides. ^{125}I -A β_{1-43} probe was incubated with hybridoma secretions from the indicated clones. A standard polyethylene glycol separation method was used to detect ^{125}I -A β_{1-43} bound antibody (Table 2). Results presented in Table 2 indicate that the antibodies generated to the peptide fragments also bound full length A β_{1-43} .

Table 2 ^{125}I -A β_{1-43} Binding Assay

Addition		^{125}I -A β_{1-43} Bound (cpm)
Control Hy		3,171
Control Hy		2,903
6E2		15,938
6E2	1/10	9,379
3B1		12,078
3B1	1/10	3,353
8E3		10,789
8E3	1/10	3,249

It was reported that when ^{125}I -A β_{1-40} is added to human plasma, ~89% binds to albumin (Biere et al., *J. of Biol. Chem.* 271(51): 32916 (1996)). This raises the concern that the bound albumin will interfere with antibody binding. Binding assays were performed in the presence and absence of serum albumin, to determine whether albumin binding interferes with antibody binding to A β . The ability of purified 5A11 monoclonal anti-A β antibody to bind ^{125}I -A β_{1-40} was unaffected by the presence of human serum albumin (HSA) at 60 mg/ml, even though this was a 500-fold molar excess over the antibody concentration

(Table 3). These results indicate that the ability of antibodies to bind to and sequester $A\beta$ in the blood will not be attenuated by the presence of other binding proteins.

Table 3. ^{125}I - $A\beta_{1-40}$ Binding to Antibody in the Presence of Human Serum Albumin*

Bound	^{125}I - $A\beta_{1-40}$ Bound	Specifically
<u>Addition</u>	<u>(cpm)</u>	<u>(% of total added)</u>
Control	8,560	-
+ 5A11 anti- $A\beta$	64,589	79
Control + HSA*	3,102	-
+ 5A11 anti- $A\beta$ + HSA*	55,304	75

*HSA at 60 mg/ml (~1 mM); anti- $A\beta$ 5A11 at 2×10^{-6} M; Added ~70,000 cpm of ^{125}I - $A\beta_{1-40}$

Monoclonal Antibody Production

A mouse was immunized with a KLH conjugate of the central region $A\beta_{10-25}$ peptide (This peptide antigen had a phenylalanine statine transition state analog at an amide linkage, discussed further in Section II, below). A hybridoma fusion was performed and the resulting monoclonal antibodies analyzed to characterize the specificity of the immune response to the vaccine. Hybridoma supernatants produced in the fusion were screened using ELISA to assess their binding to the $A\beta_{1-43}$ peptide.

The monoclonal antibodies produced were determined to bind to the $A\beta_{1-43}$ peptide adsorbed directly onto an ELISA plate. Strong color reactions were obtained in this ELISA using only 10 μ l of hybridoma supernatant while the addition of media alone produced low background color. These results indicate that the antibodies not

only bound to the small peptide immunogen but they were also reactive with the full-length $A\beta_{1-43}$. Importantly, antibodies bound to the carrier-free $A\beta$ peptide adsorbed directly onto microtitre plates, showing their
5 specificity for the peptide rather than the immunogenic carrier. The high affinity 5A11 monoclonal antibody (Table 3) was obtained from this hybridoma fusion.

A second mouse was immunized with a KLH conjugate of the $A\beta_{35-43}$ analog encompassing the C-terminal region of
10 $A\beta$. Serum from the mouse was screened for reaction with $A\beta_{1-43}$ adsorbed directly onto the ELISA wells. The assay results are presented in Table 4. The spleen of this mouse was then used for a hybridoma fusion to further characterize the specificity of its immune response.
15 Importantly, none of the mice immunized with $A\beta$ vaccines or the anti- $A\beta$ ascites-producing mice displayed ill effects even though some of those induced antibodies cross-react with mouse $A\beta$ and mouse amyloid precursor protein.

Table 4 ELISA for Binding of Antiserum Directed to the Carboxy-terminal $A\beta$ Peptide

Antibody Bound (O.D. 450 nm)	
Addition	Native $A\beta_{1-43}$
Control Serum	0.484
Mouse Antiserum	1.765

Monoclonal antibodies from hybridoma clones generated above were screened for binding to the small carboxy-terminal peptide $A\beta_{35-43}$ and the full-length $A\beta_{1-43}$. Results are presented in Figure 5. The monoclonal
5 antibodies bound to the carboxy-terminal locus on each of these carrier-free $A\beta$ peptides adsorbed directly to the microtitre plate, confirming their specificity for the peptide rather than the immunogenic carrier. The clones

were also tested with $A\beta_{1-40}$ to identify antibodies which do not react with this shortened, 40 amino acid residue version of $A\beta$ and thus will specifically bind to the carboxy-terminus of $A\beta_{1-43}$ (Fig. 5). Used therapeutically, this vaccine should elicit antibodies which will preferentially bind the less abundant, but more noxious $A\beta_{1-43}$ species in the blood as opposed to the smaller and less detrimental $A\beta_{1-40}$.

In a separate experiment, mice were immunized with a vaccine comprised of a cocktail of the three distinct KLH-peptide antigens (Figs. 2-4) representing the distinct regions of β -amyloid (Fig. 1). Control mice were immunized with KLH alone. The antigens were emulsified in complete Freund's adjuvant prior to the first injection and in incomplete Freund's adjuvant for subsequent injections. Tests were performed on diluted serum from these $A\beta$ -KLH immunized mice to determine the presence of specific anti- $A\beta$ antibodies. The $A\beta_{1-16}$, $A\beta_{14-25}$, $A\beta_{34-43}$, $A\beta_{1-40}$, and $A\beta_{1-43}$ peptides were used to identify antibody specificity. The peptides were adsorbed directly onto an ELISA plate. The results are presented in Table 5. The results indicate that mice immunized with the cocktail of the three peptide antigens produced serum containing antibodies which react with the amino-terminal, central region, and carboxyl-terminal peptides, as well as with the full-length $A\beta$ 40-mer and 43-mer. The constant presence of this spectrum of anti- $A\beta$ antibodies will be very effective in binding all of the soluble $A\beta$ in the peripheral circulation of a vaccinated animal.

Table 5 ELISA to Measure the Serum Antibodies Present in Immunized Mice

Immunogen	ELISA READING (O.D. 450 nm)				
	$A\beta_{1-16}$	$A\beta_{14-25}$	$A\beta_{14-43}$	$A\beta_{1-40}$	$A\beta_{1-43}$
Mouse 1 (Control) KLH	0.076	0.038	0.064	0.042	0.066
Mouse 2 $A\beta$ -KLH Cocktail	3.013	1.258	3.191	2.337	2.598
Mouse 3 $A\beta$ -KLH Cocktail	1.484	1.180	2.068	1.758	1.680
Mouse 4 $A\beta$ -KLH Cocktail	1.486	1.072	2.276	1.444	1.709

Vaccine Trials in Non-human Primates

Given the potential importance of β -amyloid vaccine therapy for human patients of Alzheimer's disease, a human-compatible, alum-based $A\beta$ peptide vaccine preparation has been tested in non-human primates. Antibody production and safety studies for the human-compatible β -amyloid vaccines have commenced in Cynomolgus monkeys (*Macaca fascicularis*). This animal system is highly relevant to human applications since the predicted amino acid sequence of β -amyloid in these primates is identical to humans, and their basic physiology and immunological systems closely approximate those which will be encountered in a clinical situation. Cynomolgus monkeys were vaccinated monthly and were periodically bled to monitor anti- $A\beta$ levels in the serum. The monkeys were also observed for any ill effects.

The Cynomolgus monkeys mounted a strong immune response to a single injection of the simplest vaccine preparation composed of the full length β -amyloid peptide adsorbed to an aluminum hydroxide gel. The specificity of those early anti- β -amyloid antibodies was characterized by ELISA using various $A\beta$ peptide fragments (Table 6). This analysis indicated that the monkeys produced antibodies that bind to the full-length peptide and react with its amino-terminal, central and carboxyl-terminal regions.

Table 6 ELISA to Measure the Serum Antibodies Present in $A\beta$ Vaccinated *Macaca fascicularis*

Vaccination Schedule	ELISA READING (O.D. 450 nm)				
	$A\beta_{1-16}$	$A\beta_{14-25}$	$A\beta_{34-43}$	$A\beta_{1-40}$	$A\beta_{1-43}$
Pre-Vaccination	0.511	0.404	0.370	0.380	0.235
$A\beta$ /Alum (1st month)	2.115	1.687	0.671	2.393	2.479

Importantly, the vaccinated monkeys are perfectly healthy and appear compatible with the anti- $A\beta$ antibodies that have been circulating in their body for over three months. Thus far, there are no apparent side effects due to cross-reaction of the anti- $A\beta$ antibodies with naturally occurring β -amyloid precursor protein or other vital components. These animals were closely observed by a veterinarian, and have exhibited no signs of autoimmune disease, immune complex disease or any other adverse/toxic reaction to the vaccination.

In continuing experiments boost injections will be performed as per usual methods. The sera produced will be monitored for antibody specificity and affinity parameters as the immune response intensifies and matures. At termination, a complete necropsy and histopathological examination will be performed on the monkeys. Genetically engineered $A\beta$ vaccines, discussed below, will also be evaluated in the *Cynomolgus* monkeys to determine if they will prove to be even better immunogens.

Antibodies Affect the Distribution of ^{125}I - $A\beta$ in Normal Mice

Anti- $A\beta$ antibodies in the circulation cannot cross the blood-brain barrier to a significant extent and therefore should act as a sink that prevents ^{125}I - $A\beta_{1-40}$ from reaching the brain. This retention effect was demonstrated by measuring the blood levels in mice 4 h after injecting them with equal amounts of ^{125}I - $A\beta_{1-40}$

either alone or along with our 5A11 anti-A β monoclonal antibody (Table 7). The passage of ^{125}I -A β_{1-40} out of the peripheral circulation was greatly curtailed in animals which concomitantly received the specific anti-A β antibody. That finding extends the *in vitro* results obtained with the 5A11 antibody (Table 3) by demonstrating the antibody can effectively bind A β in an experimental animal. The observation that animals treated with this antibody retained 10-times more ^{125}I -A β_{1-40} in the circulation indicates that the equilibrium distribution of A β in the body can be dramatically altered by selective sequestration in the blood.

Table 7 Anti-A β Antibody Impedes the Passage of ^{125}I -A β_{1-40} Out of the Circulation

Mouse Injected With;	^{125}I -A β_{1-40} in Blood (cpm/gm)
^{125}I -A β_{1-40} alone	27,300
^{125}I -A β_{1-40} + 5A11 anti-A β	278,900

Genetically Engineered Vaccines

Genetically engineered β -amyloid antigen vaccines for use in humans are currently being developed in order to induce protective levels of anti- β -amyloid antibodies. β -amyloid fragments will be engineered into chimeric A β vaccines which incorporate highly immunogenic carrier moieties to increase the appropriate antigenic response in a human patient. Carrier moieties suitable for use include diphtheria toxoid (DT) and the hepatitis B core antigen (HBcAg). These represent powerful delivery systems for β -amyloid peptides, and are known to induce an excellent, high titer immune response when used with alum as an adjuvant.

DT is licensed for use as a conjugate vaccine for *H. influenzae* type B and renders this immunogen T-cell dependent. The expression of DT in recombinant *E. coli*

is high. One or more of the above described β -amyloid peptides will be fused at the C-terminus of the catalytic domain of DT, or at either end of the combined transmembrane/receptor binding domains of DT. The
5 produced fusions will be used with an aluminum hydroxide gel adjuvant to generate potent vaccines.

High titers of antibody directed against heterologous epitopes have been produced using the HBcAg delivery systems and aluminum hydroxide gel adjuvant.
10 HBcAg has several distinct advantages as a fusion partner for A β peptides. The immunodominant internal site between amino acids 75 and 81 can accommodate heterologous sequences up to 45 amino acids. The core self-assembles into larger 27nm particles that are highly
15 immunogenic. Furthermore, HBcAg can be produced in recombinant *E. coli* at elevated levels.

The genetically engineered vaccines produced will be tested for effectiveness in depleting or preventing plaques using mouse and other relevant animal models.
20 Antibody production and safety trials for the vaccines will be conducted in Cynomolgus monkeys.

Methods of the Invention

Peptide synthesis. The 40mer A β_{1-40} , the 43mer A β_{1-43} , and the three small A β peptides A β_{1-16} , A β_{10-25} , and A β_{35-43} , were
25 synthesized by standard automated Fmoc chemistry. Newly synthesized peptides were purified by HPLC and their composition was verified by mass spectral and amino acid analysis. The A β 43mer was obtained from a commercial source (Bachem, Torrance, CA).

30 Conjugation of β -Amyloid Peptides to Immunogenic Carriers. The small A β peptides were linked to the KLH carrier protein in order to render them antigenic. A Cys residue was strategically placed at the N- or C-terminal end of these A β peptides to provide a suitable linkage

group for coupling them via a thioether bond to maleimide activated carrier proteins. This linkage is stable and attaches the peptide in a defined orientation. Addition of ~20 peptides/KLH is typically obtained by this conjugation method. The longer, full length A β peptides were linked to carrier proteins using a glutaraldehyde coupling procedure.

β -amyloid Antigen Cocktail. The three A β peptides shown in Figs. 2-4 were each individually conjugated to KLH. 20 μ g of each of these three conjugates was then mixed together. This mixture was emulsified with complete Freund's adjuvant and injected i.p. into mice. Subsequent monthly i.p. booster injections used the same cocktail mixture emulsified in incomplete Freund's adjuvant. Control mice received a similar immunization protocol but using KLH which had not been conjugated with the A β peptides.

Immunization of Mice. Normal BALB/c mice were immunized by standard procedures with the KLH-linked A β vaccines described above. Briefly, mice were injected i.p. with antigen emulsified in complete Freund's adjuvant, followed by a second course in incomplete Freund's adjuvant. The mice were i.v. boosted with antigen in PBS three days prior to bleeding them or removing the spleen for hybridoma fusions to produce monoclonal antibodies.

None of the mice immunized with A β vaccines or the anti-A β ascites-producing mice displayed ill effects even though some of the antibodies cross-reacted with mouse A β and mouse amyloid precursor protein.

ELISA. The presence of bound anti-peptide antibodies was revealed by using a peroxidase-labeled anti-mouse IgG probe followed by the chromogenic substrate (Engvall et al., *Immunochemistry* 8: 871-875 (1971)).

Binding Assay. Both A β ₁₋₄₃ and A β ₁₋₄₀ were radiolabeled with ¹²⁵I. The iodinated peptide was separated from unlabeled material by HPLC to give essentially quantitative specific activity (~2000 Ci/mmol) (Maggio et al., *Proc. Natl. Acad. Sci.* 89: 5462 (1992)). ¹²⁵I-A β ₁₋₄₃ probe was incubated for 1h at 23°C with Hy media taken from hybridoma clones producing monoclonal anti-A β antibodies. A standard polyethylene glycol separation method was used to detect the amount of ¹²⁵I-A β ₁₋₄₃ bound to antibody.

10 β -Amyloid Vaccines for Primates. The immunogen used was a sytheticA β peptide encompassing amino acids 1-41 of the A β protein. This peptide was purified by HPLC and freeze-dried and then resuspended in sterile water at a concentration of 1.5 mg/ml. The vaccine was prepared by
15 mixing 7.5 ml of a 2% aluminum hydroxide gel adjuvant (Alhydrogel, Superfos Biosector, Denmark), referred to herein as alum gel, with 7.5 ml of the peptide. Tests showed that all of the peptide was adsorbed to the alum gel after mixing for 12 hours at 25°C.

20 Monkeys were initially vaccinated by intramuscular (i.m.) injection of 0.5 ml of the alum-adsorbed peptide. A second vaccination (boost) of the same vaccine preparation (0.5 ml) was administered a month later. Subsequent identical monthly injections (boosts) will be
25 given until the experiment is terminated.

Genetically Engineered Vaccines. Highly immunogenic carrier moieties will be used to construct chimeric A β vaccines. Moieties used will include diphtheria toxoid (DT) and the hepatitis B core antigen (HBcAg). The HBcAg
30 expression system will be utilized (Schodel et al., *Infect. and Immun.* 57: 1347-1350 (1989); Schodel et al., *J. of Exper. Med.* 180: 1037-1046 (1994); Schodel et al., *J. of Virology* 66: 106-114 (1992); Milich et al., *Annals New York Academy of Sciences*: 187-201 (1993)). The amino
35 terminal end of the catalytic domain of HBcAg has a

signal sequence which should allow the A β fusion protein to be secreted into the culture medium. The culture medium will be concentrated using a large Amicon ultrafiltration device, and the concentrate then chromatographed on a large Superdex 75 column. Recombinant products obtained from within lysed cells will be separated from bacterial protein using a combination of anion exchange and size exclusion FPLC.

10 SECTION II: ELICITING MONOCLONAL ANTIBODIES WITH
TRANSITION STATE ANTIGENS

Transition state peptide antigens

Different types of transition state peptide antigens were synthesized to use in the generation of antibodies which preferentially recognize (hydrolysis) transition states of A β at a predetermined amide linkage position.

A series of statine (Sta) transition state analogs encompassing the carboxy-terminal region of A β (Cys-Met-Val-Gly-Gly-Val/Sta-Val/Sta-Ile/Sta-Ala-Thr) were synthesized. Replacement of the proposed scissile peptide linkage between Val₃₉ and Val₄₀, Val₄₀ and Ile₄₁, and Ile₄₁ and Ala₄₂, with a "statyl" moiety (-CHOH-CH₂-CO-NH-) was designed to elicit catalytic antibodies that hydrolytically cleave A β at one of these sites (Fig. 6). A Cys residue was placed at the N-terminal position of these peptides to provide a suitable linkage group for coupling to a maleimide-activated carrier protein.

A series of phenylalanine statine (PhSta) transition state analogs encompassing the central region of A β (Cys-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe/PhSta-Phe/PhSta-Ala-Glu-Asp-Val-Gly-amide) was synthesized in this laboratory.

Replacement of the proposed scissile peptide linkage between Phe₁₉ and Phe₂₀, and between Phe₂₀ and Ala₂₁, with a statyl moiety (-CHOH-CH₂-CO-NH-) was designed to elicit catalytic antibodies that hydrolytically cleave A β at

these sites (Fig. 7). A Cys residue was placed at the C-terminus of these peptides to provide a sulfhydryl linkage group for coupling the peptides to antigenic, maleimide-activated carrier proteins such as KLH.

5 A structural comparison (Fig. 8) was made between the native A β peptide and the transition state phenylalanine statine A β peptide using a graphics workstation. An energy minimization algorithm (2000 iterations) was applied to arrange each peptide in its
10 most favorable conformation.

 The peptide link (-CO-NH-) between Phe₁₉ and Phe₂₀ was replaced with an elongated "statyl" moiety (-CHOH-CH₂-CO-NH-) and an energy minimization was applied. This orientation shows the difference between the planar
15 peptide link (-CO-NH-) of natural A β (left) versus the extended, tetrahedral "statyl" moiety (-CHOH-CH₂-CO-NH-) in the transition state peptide (right).

 An antibody combining site complementary to a tetrahedral statine transition state analog will force
20 the planar peptide bond of the A β substrate into a transition state-like conformation. Such distortion should catalyze the cleavage of A β at that locus in the peptide sequence.

 The possibility of using a reduced peptide bond
25 linkage to mimic the transition state during hydrolysis of an amide linkage was also explored. A reduced peptide bond linkage can be easily placed at almost any site in the A β molecule to produce a reduced peptide bond transition state analog. This analog can also be used to
30 elicit catalytic antibodies that will hydrolytically cleave A β at the chosen site. The reduced peptide bond transition state A β analog made was the (Gln-Lys-Leu-Val-Phe-CH₂-NH₂⁺-Phe-Ala-Glu-Asp-Val-Gly-Cys-amide) central region peptide; [calculated 1,342 (M+1); observed 1,344].

35 A structural comparison (Fig. 9) was made between the native A β peptide and the reduced peptide bond transition state A β analog using a graphics workstation.

The peptide link (-CO-NH-) between Phe₁₉ and Phe₂₀ was replaced with a reduced peptide bond (-CH₂-NH₂⁺-) and an energy minimization was applied. The orientation shown indicates the difference between the planar peptide link (-CO-NH-) of natural A β (left) versus the corresponding tetrahedral moiety (-CH₂-NH₂⁺-) in the reduced peptide bond transition state analog (right). An energy minimization algorithm (2000 iterations) was applied to arrange each peptide in its most favorable conformation.

10 A phosphoramidate transition state analog of the carboxy-terminal region of A β has also been synthesized (Fig. 10). Replacement of the proposed scissile peptide linkage between Gly₃₈ and Val₃₉ with a phosphoramidate moiety (-PO₂⁻-NH-) was designed to elicit catalytic antibodies that will hydrolytically cleave A β at this site. The N-acetyl-Cys residue was placed at the position of Leu₃₄ to provide a suitable linkage group for coupling this peptide to an antigenic carrier protein. The structures in Figure 11 represent the putative transition state for peptide hydrolysis by zinc peptidases, versus structure of and the phosphonate and phosphoramidate mimics. Similar tetrahedral transition state intermediates are known to be formed by reaction with each of the four classes of proteolytic enzymes, the serine-, cysteine-, aspartic- and metallo-peptidases.

25 A structural comparison was made between the native A β peptide and the transition state phosphoramidate A β peptide (Fig. 12) using a graphics workstation. The peptide link (-CO-NH-) between Gly₃₈ and Val₃₉ was replaced with a phosphoramidate bond (-PO₂⁻-NH-) and an energy minimization was applied. The orientation shown in Figure 12 illustrates the difference between the planar peptide link (-CO-NH-) of native A β (left) versus the corresponding tetrahedral phosphoramidate bond (-PO₂⁻-NH-) in the transition state peptide (right).

35 An antibody combining site complementary to the tetrahedral transition state analog on the right of Fig.

-38-

12, will force the normally planar bond of the A β substrate peptide on the left into a transition state-like conformation. Such bond distortion is expected to catalyze the hydrolytic cleavage of the A β peptide at the Gly₃₈-Val₃₉ linkage.

Immunization with Transition State Peptide Antigens

Peptide antigens were coupled to the immunogenic carrier KLH prior to immunization of mice. Standard protocols were used to immunize BALB/c mice with the KLH-linked A β peptides described in the preceding sections. Briefly this procedure used i.p. injection of the different antigens emulsified in complete Freund's adjuvant, followed by a second course in incomplete Freund's adjuvant. Three days prior to hybridoma fusion, the BALB/c mice were boosted i.v. with antigen in PBS.

A hybridoma fusion was performed using the spleen of a mouse immunized with either a mixture of the phenylalanine statine transition state antigens generated (Fig. 7), a mixture of the statine (Sta) transition state A β antigens generated (Fig. 6), the reduced peptide bond transition state A β antigen generated (transition state mimic located between Phe₁₉-Phe₂₀), or the phosphoramidate transition state A β antigen generated (transition state mimic located between Gly₃₈-Val₃₉). Monoclonal antibodies listed in Table 8 were generated from these mice.

Table 8.

Analog Used	Bonds Modified	Potential Cleavage Sites	Antibodies Generated
statine	Val ₃₉ -Val ₄₀ Val ₄₀ -Ile ₄₁ Ile ₄₁ -Ala ₄₂	Val ₃₉ -Val ₄₀ Val ₄₀ -Ile ₄₁ Ile ₄₁ -Ala ₄₂	2B2, 2H6, 3F2, 4D3, 6A6, 1E4, 11E9, 9D6, 5C7, 7C7, 1D12
phenylalanine-statine	Phe ₁₉ -Phe ₂₀ Phe ₂₀ -Ala ₂₁	Phe ₁₉ -Phe ₂₀ Phe ₂₀ -Ala ₂₁	6E2, 5A11, 6F11, 2E3, 8E3, 5G4, 4C7, 8D12, 2C12, 4G7, 5C7, 3C1, 4H9, 8E6, 1H2, 3B1, 2H11
reduced peptide	Phe ₁₉ -Phe ₂₀	Phe ₁₉ -Phe ₂₀	6E7, 6F6

-39-

bond

phospho-
amidateGlu₃₈-Val₃₉Gly₃₈-Val₃₉

in progress

Demonstration of A β Binding by Generated Antibodies

It was very important to demonstrate that the anti-A β and anti-transition state A β monoclonal antibodies bound to the natural A β ₁₋₄₃ peptide which they were
 5 designed to sequester or cleave. To do this, A β ₁₋₄₀ and A β ₁₋₄₃ were radiolabeled with ¹²⁵I and the iodinated peptide was then separated from unlabeled material by HPLC. Probe was incubated with either purified anti-A β antibodies or media taken from hybridoma clones producing
 10 anti-A β antibodies. The amount of ¹²⁵I-A β ₁₋₄₃ bound to antibody was determined using a polyethylene glycol separation method. Results of the experiment are presented in Table 3.

The data in Table 3 demonstrate the ability of the
 15 purified 5A11 monoclonal anti-A β antibody to bind a high percent of ¹²⁵I-A β ₁₋₄₀. This binding assay was used to screen clones and purified antibodies (Table 3) for their ability to bind A β . Similar procedures can also serve as the basis for a competitive displacement assay to measure
 20 the relative binding strength of different unlabeled A β peptides. (Note: with very efficient catalytic antibodies this binding assay may have to be performed on ice to ensure that no cleavage of A β occurs during the 1h incubation time.) The assay rapidly identified clones
 25 producing high affinity anti-A β antibodies.

Monoclonal antibodies from hybridomas obtained using the phenylalanine statine transition state A β -KLH antigen were screened by ELISA to assess their binding to both the normal A β ₁₋₄₃ peptide and to the phenylalanine statine
 30 transition state A β peptide. Two major patterns were found (Fig. 13).

One group of antibodies (the left portion of Fig. 13) bound to the immunizing transition state peptide and cross-reacted strongly with the native $A\beta_{1-43}$ peptide (each was adsorbed directly onto the ELISA plate). The second group (the right portion) showed a high binding preference for the phenylalanine statine transition state $A\beta$ peptide and reacted minimally with native $A\beta_{1-43}$.

Strong color reactions were obtained in this ELISA using only 10 μ l of hybridoma supernatant while Hy media alone or PBS gave a low background (Fig. 13). These results demonstrate that the comparative ELISA screen, although only a semi-quantitative measure of binding, provides a means for identifying monoclonal antibodies that are highly selective for, and most reactive with, the transition state. Importantly, the experiment was performed with carrier-free $A\beta$ peptides adsorbed directly onto microtitre plates, indicating antibody specificity for $A\beta$ peptide rather than carrier.

These findings indicate that several of the generated anti- $A\beta$ transition state antibodies were unique. They bound to both the phenylalanine statine- and normal- $A\beta$ peptides. Their selective recognition of the transition state and weaker cross-reaction with native $A\beta_{1-43}$ however indicates that this binding interaction is very different from that shown by conventional anti-native $A\beta$ antibodies. It further indicates that these new antibodies may be able to force the native $A\beta$ peptide into a conformation resembling the transition state for hydrolytic cleavage. Importantly, some of the antibodies which showed only minimal binding to $A\beta_{1-43}$ in this ELISA, did display cross-reactivity with the natural peptide using a highly sensitive ^{125}I - $A\beta_{1-43}$ binding assay (Table 3).

ELISAs were also performed to investigate the binding of anti-statine analog antibodies to both the normal $A\beta_{1-43}$ peptide and to the statine transition state

-41-

A β peptide (Fig. 14). The antibodies bound to the C-terminal locus on these carrier-free A β peptides (adsorbed directly to the microtitre plate) confirming their anti-peptide specificity. Most of the antibodies
5 preferentially recognized the statine A β transition state, but cross-reacted with native A β ₁₋₄₃. This indicates that these new antibodies are able to force the native A β peptide into a conformation resembling the transition state for hydrolytic cleavage of its C-
10 terminal amino acids. Such cleavage is predicted to convert A β ₁₋₄₃ into potentially less harmful shorter peptides, like A β ₁₋₄₀ or A β ₁₋₃₉.

Clone 11E9 had the strongest preference for the statine analog and may be the most likely to have
15 catalytic activity (Fig. 14). Several clones displayed no difference in their reactivity with the native versus statine transition state A β peptide. The clones were also tested with A β ₁₋₄₀ to identify antibodies which do not react with this shortened, 40 amino acid version of A β
20 (Fig. 14). Used therapeutically, such antibodies should preferentially bind/cleave the less abundant, but more noxious A β ₁₋₄₃ species in the blood as opposed to the smaller and less detrimental A β ₁₋₄₀.

Solid Phase and TLC A β Proteolytic Assays

25 A solid phase ¹²⁵I-labeled A β assay was developed to screen anti-transition state antibody hybridoma supernatants for specific proteolytic activity. The peptide Cys-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Tyr-amide (SEQ ID NO: 5) which encompasses amino
30 acids 14-25 of A β was radiolabeled and coupled to a thiol-reactive, iodoacetyl-Sepharose gel to form an irreversible linkage. The product was incubated with anti-transition state antibody and assayed for the progressive release of soluble ¹²⁵I-peptide from the solid
35 phase matrix. Release of radioactivity from the ¹²⁵I-A β -

Sepharose was used to identify catalytic activity (Fig. 15). The assay was verified by the ability of several different proteases to rapidly hydrolyze this Sepharose-linked A β substrate. The peptide was readily accessible
5 to proteolytic cleavage as revealed by a release of soluble ^{125}I -peptide that increased with incubation time.

The results presented in Figure 15 indicate that the antibody-containing media of several clones released ^{125}I -peptide at a greater rate than other clones from this
10 fusion or the PBS and Hy medium controls. Large amounts of these antibodies can be obtained, purified and tested at higher concentrations to achieve much faster rates of cleavage and to verify that the antibodies are acting in a catalytic mode using conventional enzyme kinetics. By
15 changing the composition of the ^{125}I -peptide this same strategy can be used to assay antibodies reactive with different regions of A β .

A thin layer chromatography-based autoradiography assay was devised to obtain more definitive evidence for
20 antibody-mediated cleavage of A β . Selected anti-phenylalanine statine A β transition state clones were expanded and ascites production induced. The different monoclonal antibodies were isolated using protein A-Sepharose. Two ^{125}I -labeled peptides, A β_{1-40} and a 17-mer,
25 encompassing amino acids 9-25 of A β , were used to test for peptide cleavage. The antibodies were added to the ^{125}I -peptides, allowed to incubate and the reaction mix spotted onto polyamide thin layer sheets which were then developed in different solvents. The migration of ^{125}I -
30 products was followed by exposing the sheet using a quantitative phosphoimager system. Quantitation of the different labeled peptide fragments produced indicated that addition of the antibodies to the A β peptides lead to significant break down of the A β peptides compared to
35 the untreated peptides (PBS).

Disaggregation of β -amyloid by Monoclonal Antibodies

The self-aggregation of synthetic $A\beta$ peptides has been shown previously to lead to microscopic structures resembling amyloid plaques in the brain (Solomon et al.,
5 *Proc. Natl. Acad. Sci. USA* 94: 4109-12 (1997); Solomon et al., *Proc. Natl. Acad. Sci. USA* 93: 452-5 (1996)) which exhibit the same bright green fluorescence upon exposure to thioflavin T. These aggregates are very stable and usually require harsh detergents or strong acids to
10 dissolve. However, it has been demonstrated that the binding of certain anti- $A\beta$ monoclonal antibodies can effectively inhibit the initial aggregation of this peptide and also disaggregate preformed $A\beta$ complexes (Solomon et al., *Proc. Natl. Acad. Sci. USA* 94: 4109-12
15 (1997); Solomon et al., *Proc. Natl. Acad. Sci. USA* 93: 452-5 (1996)).

A radioactive assay was used to quickly screen the different monoclonal antibodies produced by the present experiments for an ability to dissolve preformed $A\beta$
20 aggregates, made with ^{125}I -labeled and unlabeled soluble $A\beta$ peptide. An aliquot of the labeled aggregate was incubated with either PBS, the 5A11 anti- $A\beta$ antibody or an equal amount of an irrelevant mouse antibody (7D3, anti-human transferrin receptor), and the level of
25 released radioactivity was subsequently measured (Table 9). The $A\beta$ -specific 5A11 antibody solubilized 80% of the $A\beta$ aggregates while an equal amount of the control antibody had only a minor effect, suggesting that the equilibrium was displaced by antibody-mediated binding of
30 soluble $A\beta$.

Table 9 Solubilization of $^{125}\text{I-A}\beta_{1-40}$ Aggregate by Monoclonal Anti-A β Antibody

Addition	$^{125}\text{I-A}\beta_{1-40}$ in Ppt. (cpm)	Amount Solubilized (% of PBS Control)
PBS control	3,420	-
+ 5A11 anti-A β	676	80
+ 7D3 anti-TfR	2,458	27

Production of Vectorized Anti-A β /Anti-Receptor Bispecific Antibodies

Anti-A β antibodies were linked to anti-transferrin receptor antibodies (anti-TfR) which served as vectors for delivery of the anti-A β antibodies into the brain. The 7D3 mouse monoclonal antibody was used as the anti-TfR part of the construct. 7D3 is specific for the human receptor and selectively immunostains cortical capillaries in normal human brain tissue (Recht et al., J. Neurosurg. 72: 941-945 (1990)). Antibody attachment to the receptor is not blocked by an excess of human transferrin. The epitope recognized by this antibody is therefore distant from the receptor-ligand binding site. Bispecific antibodies constructed with this 7D3 antibody and an anti-A β antibody are predicted to be useful for therapy in patients with Alzheimer's disease.

For studies in mouse models of Alzheimer's disease an anti-mouse transferrin receptor monoclonal antibody produced in the rat was obtained. This antibody also appears to recognize a transferrin receptor epitope which does not involve ligand binding. The antibody therefore has no effect on cell proliferation when using murine lines.

A series of functional assays were performed after completion of the synthesis, purification and size analysis of the anti-A β /anti-transferrin receptor

bispecific antibodies. The vectorized bispecific antibody, composed of a rat monoclonal antibody directed against the mouse transferrin receptor plus the 5A11 mouse anti-A β monoclonal antibody, was tested for the ability to attach to transferrin receptor bearing mouse cells. Both components of the bispecific antibody were detected on the cell membrane by cytofluorimetry (Fig. 16) when this duplex was reacted with transferrin receptor positive mouse cells and probed using either a rat IgG-specific or mouse IgG-specific fluorescent secondary antibody reagent.

The capacity of the hybrid reagent to bind ^{125}I -A β compared favorably with that of the parent anti-A β antibody (Table 10).

Table 10 ^{125}I -A β Binding to Bispecific Antibody

Addition	^{125}I -A β_{1-40} Bound
	(cpm)
Control	4,199
+ anti-A β	23,301
+ anti-A β /anti-receptor	22,850

To ensure that both of these binding activities resided on the bispecific antibody, transferrin receptor positive cells were treated with the hybrid reagent, unbound material was washed away, and then the cells with bound antibody was exposed to ^{125}I -A β_{1-40} . After washing away unbound A β , the cell-bound radioactivity was compared to control cells which had been identically prepared except for omission of pretreatment with bispecific antibody. The results are presented in Table 11, and verify the dual specificity of this bispecific antibody by clearly showing that it can simultaneously attach to the cell membrane and bind ^{125}I -A β_{1-40} .

Table 11 Bispecific Antibody-Mediated Binding of ^{125}I -A β to Receptor-Positive Cells

<u>Pretreatment of Cells</u>	<u>^{125}I-Aβ_{1-40} Bound (cpm)</u>
None	2,367
+ anti-A β /anti-transferrin receptor	11,476

Transcytosis of Bispecific Antibody into the Brain

A rat monoclonal anti-mouse transferrin receptor antibody was coupled to a mouse monoclonal antibody (obtained from American Type Culture Collection (ATCC TIB 219), also designated R17 217.1.3 (*Cell. Immunol.* 83: 14-25 (1984)), so that the entry of this new vectorized bispecific construct into brain could be monitored. The bispecific antibody was labeled with ^{125}I and injected i.v. into normal mice. After different lengths of time the mice were sacrificed and the amount of ^{125}I -bispecific antibody that crossed the blood-brain barrier and entered the brain was gauged by a mouse capillary depletion method (Friden et al., *J. Pharm. Exper. Ther.* 278: 1491-1498 (1996); Triguero et al., *J. Neurochem.* 54: 1882-1888 (1990)).

The amount of vectorized bispecific antibody found in the brain parenchyma or brain capillary fractions was measured following differential density centrifugation of the brain homogenate. These values were plotted as a function of time after i.v. injection (Fig. 17). The time-dependent redistribution of radiolabeled bispecific antibody from the capillaries and into the parenchyma was consistent with its passage across the cerebral endothelial blood-brain barrier (Joachim et al., *Nature* 341: 6239:226-30 (1989)). Even greater accumulation in the parenchyma is expected to occur if the antibodies attach to A β in the cerebral plaques of plaque-bearing mice.

Monitoring the Brain Distribution of Bispecific Antibody
in Live Mice

The ability to follow the entry and accumulation of vectorized bispecific antibodies in the brain of live mice would greatly assist in the development of the intracerebral treatment of plaque-bearing mice. Such a development would enable time-course studies and would greatly reduce problems with inter-mouse variability. Preliminary studies with ^{125}I -labeled bispecific antibodies were performed to determine if immunoscintigraphy was feasible in this system. As a first step, either the radiolabeled vectorized bispecific antibody (^{125}I -R17/5A11) or a non-vectorized control bispecific antibody were administered to separate mice. Sequential brain images were accumulated at 1, 6, 24 and 48 hours following i.v. administration of the ^{125}I -labeled bispecific antibody probes. Although this technique suffered from a difficulty in determining how much of the signal was due to the levels of blood-borne radioactivity circulating through the brain, significant distinctions were noted in the brain of mice treated with the mouse transferrin receptor reactive bispecific antibody versus those receiving the control bispecific antibody. When the vectorized agent was used, brain levels increased between 1 and 6 hrs and then declined to a much lower level at 24 and 48 hrs. Mice treated with the control displayed no increase between 1 and 6 hrs. The reason for decreased brain levels at 24 hrs and beyond is not known but might be due to dehalogenation of the bispecific antibody probes so that free ^{125}I is released. Alternative methods utilizing radioactive labels such as ^{111}In (Sheldon et al., Nucl. Med. Biol. 18: 519-526 (1991)) or $^{99\text{m}}\text{Tc}$ (Texic et al., Nucl. Med. Biol. 22: 451-457 (1995)) attached to the vectorized bispecific antibody can be utilized in future experiments if the use of iodine presents a technical problem. This imaging technology will be useful for determining if smaller

vectorized bispecific antibodies (eg. $F(ab')_2$) with different physical properties and an altered biodistribution will penetrate into the brain more effectively.

5 $F(ab')_2$ Heterodimers for Vector-Mediated Transport into the Brain

The introduction of whole antibodies into the brain might be detrimental if they were to fix complement and promote complement-mediated lysis of neuronal cells. The
10 development of smaller vectorized $F(ab')_2$ bispecific reagents is expected to avoid this problem. It has been shown that aggregated $A\beta$ itself can fix complement in the absence of any antibody and that the resulting inflammation may contribute to the pathology of
15 Alzheimer's disease. The possibility of intracerebral antibody having a similar effect would be greatly reduced by eliminating the Fc region of the antibody. Moreover, since coupling of Fab' halves uses the intrinsic hinge region cysteines, no extraneous substituent linkage
20 groups need be added.

Faster or more efficient entry into the brain represents another potential advantage that smaller $F(ab')_2$ or Fv_2 reagents provide for intracerebral delivery. Such modified bispecific agents can be
25 prepared and compared to full-sized hybrid antibodies for their relative effectiveness in reaching the brain, crossing the blood-brain barrier, and affecting $A\beta$ plaque development, by the methods described herein. It is important to note, however, that only minor differences
30 were found when the capacity of differently-sized anti-transferrin receptor bispecific reagents for delivering toxins into cells by receptor-mediated endocytosis was compared (Raso et al., *J. Biol. Chem.* 272: 27623-27628 (1997)). This observation might indicate that little
35 variation will be seen for transcytosis across the brain capillary endothelial cells which form the blood-brain

barrier. At the very least however one would expect the two types of vectorized molecules to have different biodistribution and plasma half-life characteristics (Spiegelberg et al., *J. Exp. Med.* 121: 323 (1965)).

5 Methods of the Invention

Antigen Synthesis. The statine and phenylalanine statine transition state peptides were synthesized using automated Fmoc chemistry. Fmoc-statine (Sta), [N-Fmoc-(3S,4S)-4-amino-3-hydroxy-6-methyl heptanoic acid] and
10 Fmoc-"phenylalanine statine" (PhSta), [N-Fmoc-(3S,4S)-4-amino-3-hydroxy-5-phenylpentanoic acid] were purchased commercially. Each peptide was tested for purity by HPLC and its composition was verified by mass spectral and amino acid analysis.

15 The design strategy and methods for synthesizing phosphoramidate- and phosphonate-based transition state peptides are straightforward (Bartlett et al., *Am. Chem. Society* 22: 4618-4624 (1983); Bartlett et al., *Biochemistry* 26: 8553-8561 (1987)). The N-terminal
20 portion of the peptide (N-acetyl-Cys-Met-Val-Gly) was made using standard automated Fmoc chemistry. After cleavage from the resin the N-acetyl tetrapeptide was treated with pyridine disulfide to protect its sulfhydryl group. An acid chloride of Cbz-glycine phosphonate
25 monomethyl ester (Bartlett et al., *Am. Chem. Society* 22: 4618-4624 (1983); Bartlett et al., *Biochemistry* 26: 8553-8561 (1987)) was coupled with Val-Val-Ile-Ala-amide which was synthesized by automated Fmoc chemistry. The last
30 amino acid of A β , Thr, was omitted due to potential problems with its unprotected hydroxyl group. The product, Cbz-Gly-PO₂⁻-NH-Val-Val-Ile-Ala-amide has a phosphoramidate (methyl ester) bond between the Gly and Val residues. Next, the Cbz blocking group was removed using hydrogen so that the protected N-acetyl-Cys-Met-
35 Val-Gly peptide could be added to the amino terminal end

-50-

of this transition state peptide by HBTU-activated peptide linkage. Treatment with mercaptoethanol and rabbit liver esterase was used to deblock the peptide. Each key component in the synthetic scheme was tested for
5 purity by HPLC and its composition was verified by mass spectral and amino acid analysis.

A reduced peptide bond linkage was placed at the indicated sites in the A β molecule. Automated Fmoc chemistry was used to begin synthesis of the peptide. A
10 pre-synthesized Fmoc amino aldehyde was then added manually and after the imide was reduced, automated synthesis was resumed (Meyer et al., *J. Med. Chem.* 38: 3462-3468 (1995)).

Coupling of Antigen to Carrier. The native and
15 transition state A β peptides were coupled to maleimide-activated KLH by standard procedures (Partis et al., *J. Pro. Chem.* 2: 263-277 (1983)), in order to elicit an immune response. A Cys residue was strategically placed at the N- or C-terminal end of the peptides to provide a
20 suitable linkage group for coupling them via a thioether bond to maleimide activated carrier proteins. This stable linkage attaches the peptide in a defined orientation. Addition of ~20 peptides/KLH has been obtained based upon the transition state amino acid
25 content as determined by amino acid analysis of the hydrolyzed conjugates (Tsao et al., *Anal. Biochem.* 197: 137-142 (1991)).

Immunization of Mice. Standard protocols were used to immunize mice with the KLH-linked A β peptides described
30 in the preceding sections. Briefly this procedure used i.p. injection of the different antigens emulsified in complete Freund's adjuvant, followed by a second course in incomplete Freund's adjuvant. Three days prior to the hybridoma fusion, the BALB/c mice were boosted i.v. with
35 antigen in PBS.

-51-

After -1 month animals were given a boost i.p. using the antigen emulsified with incomplete adjuvant. Serum from these animals was analyzed for anti-peptide antibodies by ELISA. BALB/c mice showing abundant
5 antibody production were boosted by an i.v. injection with antigen and three days later they were used to generate hybridoma clones that secrete monoclonal antibodies.

None of the mice immunized with A β vaccines or the
10 anti-A β ascites-producing mice displayed ill effects even though some of those induced antibodies cross-react with mouse A β and mouse amyloid precursor protein.

Hybridoma Production I. A hybridoma fusion was performed using the spleen of a mouse immunized with the
15 phenylalanine statine transition state A β -KLH antigen. Spleen cells from mice with the highest titre were fused with mouse myeloma NS-1 cells to establish hybridomas according to standard procedures (Köhler et al., Nature 256: 495 (1975); R.H. Kennett, Fusion Protocols.
20 Monoclonal Antibodies, eds. R.H. Kennett, T.J. McKearn and K.B. Bechtol. Plenum Press, New York. 365-367 pp. (1980)).

¹²⁵I-A β Binding Assay. A β ₁₋₄₀ and A β ₁₋₄₃ were radiolabeled with ¹²⁵I and the iodinated peptide then separated from
25 unlabeled material by HPLC to give quantitative specific activity (~2000 Ci/mmol) (Maggio et al., Proc. Natl. Acad. Sci. 89: 5462-5466 (1992)). This probe was incubated for 1h at 23°C with either purified anti-A β antibodies or media taken from hybridoma clones producing
30 anti-A β antibodies. A polyethylene glycol separation method was used to detect the amount of ¹²⁵I-A β ₁₋₄₃ bound to antibody. By using serial dilution, this assay can provide relative binding affinities for the different hybridoma supernatants or purified antibodies.

Solid Phase A β Proteolytic Assay. A solid phase ^{125}I -labeled A β assay was developed to screen anti-transition state antibody hybridoma supernatants for specific proteolytic activity. The Cys-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Tyr-amide peptide (SEQ ID NO: 5) encompassing amino acids 14-25 of A β was radiolabeled with ^{125}I and the iodinated peptide was then separated from unlabeled material by HPLC. The highly radioactive A β peptide was coupled to a thiol-reactive, iodoacetyl-Sephadex gel to form an irreversible linkage. Antibodies were added to the labeled A β , which was then assayed for progressive release of soluble ^{125}I -peptide from the solid phase matrix at pH 7, 25°C. This assay was verified by the ability of several different proteases in to rapidly hydrolyze this Sephadex-linked A β substrate. Release of soluble ^{125}I -peptide increased with incubation time.

Although A β is cleaved by several naturally occurring proteases, preliminary tests indicated that interference from high levels of background hydrolysis was not a problem when assaying hybridoma supernatants of clones that did produce catalytic antibodies. A further precaution that can be taken against exogenous proteases is carrying out all hybridoma cell fusions and cell culturing in serum-free media.

TLC A β Proteolytic Assay. A thin layer chromatography-based autoradiography assay was used to obtain more definitive evidence for antibody-mediated cleavage of A β . Selected anti-phenylalanine statine A β transition state clones were expanded and ascites production induced. The different monoclonal antibodies were isolated using protein A-Sephadex. The cleavage assay used ^{125}I -A β_{1-40} and an ^{125}I -labeled 17-mer, encompassing amino acids 9-25 of A β . Binding of the two ^{125}I -labeled peptides to the purified monoclonal antibodies 5A11 and 6E2 was examined using either a PEG precipitation assay or by a co-

electrophoresis method. Peptide cleavage was tested by adding the antibodies to the ^{125}I -peptides, incubating and then spotting the reaction mix onto polyamide thin layer sheets. The chromatographs were developed in different solvents (eg. 0.5 N HCl, 0.5 N NaOH or pH 7 phosphate buffer) and the migration of ^{125}I -products was followed by exposing the sheet using a quantitative phosphoimager system.


Screen and Isolate Select Anti-A β Antibodies. An ELISA was used to initially screen for anti-A β and anti-transition state A β peptide monoclonal antibodies. Both the transition state peptide and the corresponding natural A β peptide were adsorbed onto separate microtitre plates. The hybridoma supernatants were screened using two assays so that the relative binding to both native and transition state A β peptides could be quantitated. Clones producing monoclonal antibodies that preferentially recognized the transition state or bound A β with high affinity were selected for expansion and further study.

Propagation and Purification of Monoclonal Antibodies. Selected clones producing anti-A β antibodies and clones producing anti-receptor antibodies were injected into separate pristane-primed mice. Ascites were collected and the specific monoclonal antibodies isolated. Purification of antibodies from ascites was accomplished using a Protein A column or alternatively, antibodies were isolated from ascites fluid by $(\text{NH}_4)_2\text{SO}_4$ precipitation and passage over an S-300 column to obtain the 150 kDa immunoglobulin fraction. Monovalent Fab fragments were prepared and isolated by established methods. Their purity was evaluated by SDS-PAGE under reducing and non-reducing conditions. 50-100 mg of purified monoclonal antibody was routinely obtained from each ascites-bearing mouse.

Further Characterization of Catalytic Activity on A β Substrates. To fully define the hydrolytic properties of the isolated anti-transition state antibodies some very important controls can be run. First the ability to
5 completely block catalytic antibody activity with the appropriate transition state peptide can be verified. This non-cleavable "inhibitor" should bind much more tightly to the antibody combining sites and thereby prevent substrate binding or cleavage. Substrate
10 specificity can be further established by showing no cleavage of a sham A β peptide having a different amino acid sequence. The products of hydrolysis can also be fully characterized by HPLC, amino acid and mass spectral analysis. Control antibodies that are not directed
15 against the transition state A β can be tested and confirmed to produce no catalysis. Finally, catalytic activity can be shown to reside in the purified Fab fragments of the anti-transition state antibody.

Purified Anti-A β Antibodies Dissolve Preformed A β Aggregates. (Walker et al., Soc. Neurosci. Abstr. 21: 257 (1995), Zlokovic, B.V., Life Sciences 59: 1483-1497 (1996)). A β precipitates were formed and measured in
20 vitro (Yankner et al., Science 250: 279-282 (1990), Kowall et al., Proc. Natl. Acad. Sci. 88: 7247-7251 (1991)). A radioactive assay was used to quickly screen the different monoclonal antibodies produced for an ability to dissolve preformed A β aggregates. After
25 adding ¹²⁵I-A β to unlabeled soluble peptide, aggregates were formed by bringing the solution to pH 5 or by stirring it overnight in PBS. An aliquot of the labeled
30 aggregate was incubated for 1 hr with either PBS, the 5A11 anti-A β antibody or an equal amount of an irrelevant mouse antibody (7D3, anti-human transferrin receptor). After centrifugation, the level of radioactivity in the
35 precipitate was measured.

Generation of Vectorized Anti-A β /Anti-Receptor Bispecific Antibodies. The anti-A β antibodies were chemically coupled to anti-human transferrin receptor and anti-mouse transferrin receptor antibodies by different methods
5 (Raso et al., *J. Biol. Chem.* 272: 27623-27628 (1997); Raso et al., Monoclonal antibodies as cell targeted carriers of covalently and non-covalently attached toxins. *In* Receptor mediated targeting of drugs, vol. 82. G. Gregoriadis, G. Post, J. Senior and A. Trouet,
10 editors. NATO Advanced Studies Inst., New York. 119-138 (1984)). A rapid thioether linkage technique was used to form strictly bispecific hybrids using Traut's reagent and the heterobifunctional SMBP reagent. One component was sparingly substituted with thiol groups (SH). These
15 readily reacted to form a thioether linkage upon mixture with the maleimido-substituted (M) second component following the reaction:



Gel filtration of the reaction mixture on an S-300
20 column yielded the purified dimer which was 300 kDa and had two sites for binding A β plus two sites for attachment to transferrin receptors on brain capillary endothelial cells. Non-targeted control hybrids were formed by linking a nonspecific MOPC antibody to the
25 anti-A β antibody. This hybrid antibody does bind A β , but, being non-reactive with transferrin receptors, should not cross the blood-brain barrier.

F(ab')₂ fragments of the two different antibody types can similarly be thioether-linked to form Fc-devoid
30 reagents that cannot bind complement which might otherwise cause neurotoxic effects. These smaller bispecific hybrids (100 kDa) can be formed by reducing the intrinsic disulfides which link the heavy chains of F(ab')₂ fragments (Raso et al., *J. Immunol.* 125: 2610-
35 2616 (1980)). The thiols generated are stabilized and

-56-

Ellman's reagent (E) is used to activate these groups on one of the components (Brennan et al., Science 229: 81-83 (1985)). Exclusively bispecific F(ab')₂ hybrids can be formed upon mixing the reduced Fab' with an activated Fab' having the alternate specificity according to the reaction:



Purification on an S-200 column will isolate hybrids with one site for binding A β and one site for interaction with the target epitope on the brain capillary endothelial cells.

A similar approach can be used to make even smaller disulfide-linked single chain Fv heterobispecific dimers, Fv_A-SS-Fv_B (50 kDa), to cross the blood-brain barrier. Soluble Fvs can be constructed to possess a carboxyl-terminal cysteine to facilitate the disulfide exchange shown in the reaction below, and create 50 kDa heterodimers exclusively:



In side by side comparisons between whole antibody and either Fab' or Fv based bispecific reagents, the latter have proven to be moderately more effective on a molar basis for cell uptake via the transferrin receptor-mediated pathway (Raso et al., J. Biol. Chem. 272: 27623-27628 (1997)). Since these smaller constructs are monovalent for the cell-surface epitope, those findings dispel the notion that cross-linking of two surface receptors is necessary for the cellular uptake of immunocomplexes.

Functional Assays for Dual Binding Activity of Bispecific Antibodies. The capacity of the hybrid reagent to bind ¹²⁵I-A β was compared with that of the parent anti-A β

antibody in a standard PEG binding assay (see Table 10 for binding assays).

The ability of the appropriate bispecific antibodies to attach to transferrin receptor bearing human or mouse cells was confirmed by cytofluorimetry. The bispecific antibody was reacted with transferrin receptor positive human or mouse cells and probed using either a rat IgG-specific or mouse IgG-specific fluorescent secondary antibody reagent.

10 Measurement of A β Binding Using 125 I-A β and a Polyethylene Glycol Separation. To ensure bispecificity, hybrid reagents were tested for a capacity to mediate the attachment of 125 I-A β to receptor-bearing cells. Transferrin receptor positive cells were treated with the hybrid reagent, washed away unbound material and then exposed these cells to 125 I-A β ₁₋₄₀. The cells were washed and the amount of cell-bound radioactivity was compared to control cells which had been identically prepared except that pretreatment with bispecific antibody was omitted.

20 Capillary Depletion. The bispecific antibody was labeled with 125 I and injected i.v. into normal mice. After different lengths of time the mice were sacrificed and the amount of 125 I-bispecific antibody that crossed the blood-brain barrier and entered the brain was gauged by a mouse capillary depletion method (Friden et al., *J. Pharm. Exper. Ther.* 278: 1491-1498 (1996); Triguero et al., *J. Neurochem.* 54: 1882-1888 (1990)). The amount of vectorized bispecific antibody found in the brain parenchyma or brain capillary fractions was measured following differential density centrifugation of the brain homogenate. These values were plotted as a function of time after i.v. injection. Progressive passage from capillaries into the parenchyma indicates active transcytosis across the blood-brain barrier.

- 58 -

Immunoscintigraphy. A non-invasive method for monitoring intracerebral delivery process which involves visualizing the entry of a radiolabeled bispecific antibody into the brain of live mice, can also be used. Radiolabeled

5 vectorized bispecific antibody (^{125}I -R17/5A11) or a non-vectorized control bispecific antibody were administered to separate mice. Sequential brain images were accumulated at 1, 6, 24 and 48 hours following i.v. administration of the ^{125}I -labeled bispecific antibody

10 probes. The animals were chemically immobilized during exposure using ketamine/xylazine anesthesia. This imaging technology could be very useful for determining if circulating anti-A β antibodies will prevent i.v. administered ^{125}I -A β from entering the brain. Digital

15 scintigraphy data was quantified using standards and the integration functions provided in the analysis software.

CLAIMS

1. An antibody which catalyzes hydrolysis of β -amyloid at a predetermined amide linkage.
2. The antibody of Claim 1 which catalyzes hydrolysis of the amide linkage between residues 39 and 40 of β -amyloid.
3. The antibody of Claim 1 which catalyzes hydrolysis of the amide linkage between residues 40 and 41 of β -amyloid.
4. The antibody of Claim 1 which catalyzes hydrolysis of the amide linkage between residues 41 and 42 of β -amyloid.
5. The antibody of Claim 1 which preferentially binds a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage, and also binds to natural β -amyloid with sufficient affinity to detect using an ELISA.
6. The antibody of Claim 1 which preferentially binds a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage, and does not bind natural β -amyloid with sufficient affinity to detect using an ELISA.
7. A vectorized antibody which is characterized by the ability to cross the blood brain barrier and the ability to catalyze the hydrolysis of β -amyloid at a predetermined amide linkage.

8. The vectorized antibody of Claim 7 which is a bispecific antibody.
9. The vectorized antibody of Claim 8 which has a first specificity for the transferrin receptor and a second specificity for a transition state adopted by β -amyloid during hydrolysis.
10. The vectorized antibody of Claim 9 which catalyzes hydrolysis of β -amyloid between residues 39 and 40.
11. A method for sequestering free β -amyloid in the bloodstream of an animal, comprising the steps:
 - a) providing antibodies specific for β -amyloid; and
 - b) intravenously administering the antibodies to the animal in an amount sufficient to increase retention of β -amyloid in the circulation.
12. A method for sequestering free β -amyloid in the bloodstream of an animal, comprising the steps:
 - a) providing an antigen comprised of an epitope which is present on endogenous β -amyloid; and
 - b) immunizing the animal with the antigen of step a) under conditions appropriate for the generation of antibodies which bind endogenous β -amyloid.
13. A method for reducing levels of β -amyloid in the brain of an animal, comprising the steps:
 - a) providing antibodies specific for β -amyloid endogenous to the animal; and
 - b) intravenously administering the antibodies to the animal in an amount sufficient to increase retention of β -amyloid in the circulation of the animal.

14. The method of Claim 13 wherein the antibodies specific for β -amyloid are catalytic antibodies which catalyze hydrolysis of β -amyloid at a predetermined amide linkage.
15. The method of Claim 13 wherein the antibodies are monoclonal.
16. The method of Claim 13 wherein the antibodies are polyclonal.
17. The method of Claim 13 wherein the antibodies specifically recognize epitopes on the C-terminus of β -amyloid₁₋₄₃.
18. A method for reducing levels of β -amyloid in the brain of an animal, comprising the steps:
 - a) providing an antigen comprised of an epitope which is present on β -amyloid endogenous to the animal; and
 - b) immunizing the animal with the antigen of step a) under conditions appropriate for the generation of antibodies which bind endogenous β -amyloid.
19. The method of Claim 18 wherein the antigen is a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage.
20. The method of Claim 18 wherein the antigen is comprised of $A\beta_{10-25}$.
21. The method of Claim 19 wherein the antibodies generated have a higher affinity for the transition state analog than for natural β -amyloid.

22. The method of Claim 19 wherein the antibodies generated catalyze hydrolysis of endogenous β -amyloid.
23. A method for preventing the formation of amyloid plaques in the brain of an animal, comprising the steps:
 - a) providing an antigen comprised of an epitope which is present on β -amyloid endogenous to the animal; and
 - b) immunizing the animal with the antigen of step a) under conditions appropriate for the generation of antibodies which bind endogenous β -amyloid.
24. The method of Claim 23 wherein the antigen is a transition state analog which mimics the transition state adopted by β -amyloid during hydrolysis at a predetermined amide linkage.
25. A method for reducing levels of circulating β -amyloid in an animal, comprising the steps:
 - a) providing an antigen comprised of an epitope which is a mimic of a predetermined hydrolysis transition state of a β -amyloid polypeptide endogenous to the animal; and
 - b) immunizing the animal with the antigen of step a) under conditions appropriate for the generation of antibodies to the β -amyloid hydrolysis transition state.
26. A method for reducing levels of circulating β -amyloid in an animal, comprising the steps:
 - a) providing antibodies which catalyze the hydrolysis of β -amyloid endogenous to the animal; and

-63-

- b) intravenously administering the antibodies to the animal.
27. A method for preventing the formation of amyloid plaques in the brain of an animal, comprising the steps:
- a) providing antibodies which catalyze hydrolysis of β -amyloid produced by the animal at a predetermined amide linkage; and
 - b) administering the antibodies to the animal in an amount sufficient to cause a significant reduction in β -amyloid levels in the blood of the animal.
28. A method for reducing levels of β -amyloid in the brain of an animal, comprising the steps:
- a) providing vectorized bispecific antibodies competent to transcytose across the blood brain barrier, which catalyze hydrolysis of β -amyloid of the animal at a predetermined amide linkage; and
 - b) intravenously administering the antibodies to the animal.
29. The method of Claim 28 wherein the vectorized bispecific antibodies specifically bind the transferrin receptor.
30. The method of Claim 28 wherein the vectorized bispecific antibodies catalyze hydrolysis of the amide linkage between residues 39 and 40 of β -amyloid.
31. A method for disaggregating amyloid plaques present in the brain of an animal comprising the steps:

-64-

- a) providing vectorized bispecific antibodies competent to transcytose across the blood brain barrier, which catalyze hydrolysis of β -amyloid produced by the animal at a predetermined amide linkage; and
 - b) intravenously administering the antibodies to the animal in an amount sufficient to cause significant reduction in β -amyloid levels in the brain of the animal.
32. A method for disaggregating amyloid plaques present in the brain of an animal, comprising the steps:
- a) providing antibodies which catalyze hydrolysis of β -amyloid produced by the animal at a predetermined amide linkage; and
 - b) administering the antibodies to the animal.
33. A method for generating antibodies which catalyze hydrolysis of a protein or polypeptide comprising the steps:
- a) providing an antigen, the antigen being comprised of an epitope which has a statine analog which mimics the conformation of a predetermined hydrolysis transition state of the polypeptide;
 - b) immunizing an animal with the antigen under conditions appropriate for the generation of antibodies to the hydrolysis transition state.
34. The method of Claim 33 wherein the protein is β -amyloid.
35. A method for generating antibodies which catalyze hydrolysis of a protein or polypeptide comprising the steps:

-65-

- a) providing an antigen, the antigen being comprised of an epitope which has a reduced peptide bond analog which mimics the conformation of a predetermined hydrolysis transition state of the polypeptide;
 - b) immunizing an animal with the antigen under conditions appropriate for the generation of antibodies to the hydrolysis transition state.
36. The method of Claim 35 wherein the protein is β -amyloid.

H₂N-DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA¹T-CO₂H (SEQ ID NO:1)
 1 10 20 30 40

FIG. 1

H₂N-Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-Gln-Lys-Cys-amide (SEQ ID NO:2)
 1 5 10 15

FIG. 2

1/13

Cys-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-amide (SEQ ID NO:3)
 1 10 15 20 25

FIG. 3

Cys-Met-Val-Gly-Gly-Val-Val-Ile-Ala-Thr-CO₂H (SEQ ID NO:4)
 35 40

FIG. 4

2/13

□ BINDING TO A835-43
 ■ BINDING TO A81-43
 ▨ BINDING TO A81-40




FIG. 5

3/13




FIG. 6







FIG. 7

4/13

**FIG. 8**


5/13

Val-Phe-CH₂-NH₂⁺-Phe-Ala


Val-Phe-CO-NH-Phe-Ala

FIG. 9


6/13

**FIG. 10**


7/13



Phosphonamidate




Phosphonate




Transition-state

FIG. 11

8/13



↑
Gly-Gly-PO₂-NH-Val-Val



↑
Gly-Gly-CO-NH-Val-Val

FIG. 12

9/13

□ BINDING TO PHENYLALANINE STATINE-AB TRANSITION STATE
 ■ BINDING TO AB1-43




FIG. 13

10/13






FIG. 14


11/13

**FIG. 15**

12/13

**FIG. 16**

13/13

**FIG. 17**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/16551

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 9/00
US CL : 435/188.5
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/188.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAPLUS & EAST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/06066 (S. TRASCIATTI, et al.) 11 February 1999, see entire document.	1-6
A	US 5,439,812 (BENKOVIC, et al.) 08 AUGUST 1995, see entire document.	1-10, 33-36
A,P	US 6,043,069 (KOENTGEN, et al.) 28 MARCH 2000, column 15, lines 17-32.	1-10, 33-36

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T"
"A" document defining the general state of the art which is not considered to be of particular relevance	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

12 OCTOBER 2000

Date of mailing of the international search report

17 NOV 2000

Name and mailing address of the ISA/US
Hoskins, Brian / 2110 West 30th Street, Suite 100
Box PCT

Authorized officer
Telephone No. (703) 306-0106

[Handwritten signature]

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/16551

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-10, 33-36

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

Form PCT/ISA/210 (continuation of Form PCT/ISA/210) of 10/1999

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/16551

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-10 and 33-36, drawn to a catalytic antibody which hydrolyzes β -amyloid, a vectorized antibody with the ability to cross the blood barrier and hydrolyze β -amyloid and a method for generating antibodies.

Group II, claims 11-12, drawn to a method for sequestering free β -amyloid in the bloodstream.

Group III, claims 13-22 and 27-32, drawn to a method for reducing levels of β -amyloid in the brain and disaggregating amyloid plaques in the brain.

Group IV, claims 23-24, drawn to a method for preventing the formation of amyloid plaques in the brain.

Group V, claims 25-26, drawn to a method for reducing levels of circulating β -amyloid in an animal.

The inventions listed as Groups I-V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Group I is drawn to a catalytic antibody and a method of generating antibodies. Groups II-V involve various uses for the catalytic antibody and each use is separate and distinct from the others.